

Polyclonal Antibody 产品货号: APRab05035



产品概述 (Summary)

产品名称 (Production Name) Moesin/Ezrin/Radixin (phospho Thr558) Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application) WB,IHC,ICC/IF,ELISA 种属反应性 (Reactivity) Human, Mouse, Rat

产品性能 (Performance)

Unconjugated 偶联物 (Conjugation) 修饰 (Modification) Phosphorylated

同种型 (Isotype) IgG

Polyclonal 克隆 (Clonality) 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) **MSN**

MSN; Moesin; Membrane-organizing extension spike protein; RDX; Radixin; 别名 (Alternative Names)

EZR; VIL2; Ezrin; Cytovillin; Villin-2; p81

基因 ID (Gene ID) 4478/5962

P26038/P35241/P15311.The antiserum was produced against synthesized

蛋白 ID (SwissProt ID) peptide derived from human Moesin/Ezrin/Radixin around

phosphorylation site of Thr558. AA range:524-573

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:5000-1:10000

E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技支持) Web: https://www.enkilife.cn Tel: 027-87002838







蛋白分子量 (Molecular Weight) 67kDa


研究背景 (Background)

Moesin (for membrane-organizing extension spike protein) is a member of the ERM family which includes ezrin and radixin. ERM proteins appear to function as cross-linkers between plasma membranes and actin-based cytoskeletons. Moesin is localized to filopodia and other membranous protrusions that are important for cell-cell recognition and signaling and for cell movement. [provided by RefSeq, Jul 2008], function: Probably involved in connections of major cytoskeletal structures to the plasma membrane., PTM: Phosphorylation on Thr-558 is crucial for the formation of microvilli-like structures., similarity: Contains 1 FERM domain., subcellular location: Phosphorylated form is enriched in microvilli-like structures at apical membrane, subunit: In resting T-cells, part of a PAG1-SLC9A3R1-MSN complex which is disrupted upon TCR activation (By similarity). Binds SLC9A3R1, tissue specificity: In all tissues and cultured cells studied.,


研究领域(Research Area)

Leukocyte transendothelial migration; Regulates Actin and Cytoskeleton;


图片 (Image Data)



Immunofluorescence analysis of A549 cells, using Moesin/Ezrin/Radixin (Phospho-Thr558) Antibody. The picture on the right is blocked with the phospho peptide.



E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838






Polyclonal Antibody 产品货号: APRab05035

Immunohistochemistry analysis of paraffin-embedded human pancreas, using Moesin/Ezrin/Radixin (Phospho-Thr558)

Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from NIH/3T3 cells, using Moesin/Ezrin/Radixin (Phospho-Thr558) Antibody. The lane on the right is blocked with the phospho peptide.

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838