

产品名称: MyoD (phospho Ser200) Rabbit Polyclonal Antibody
产品货号: APRab05054



产品概述 (Summary)

产品名称 (Production Name)	MyoD (phospho Ser200) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	MYOD1
别名 (Alternative Names)	MYOD1; BHLHC1; MYF3; MYOD; Myoblast determination protein 1; Class C basic helix-loop-helix protein 1; bHLHc1; Myogenic factor 3; Myf-3
基因 ID (Gene ID)	4654.0
蛋白 ID (SwissProt ID)	P15172.The antiserum was produced against synthesized peptide derived from human MYOD around the phosphorylation site of Ser200. AA range:171-220

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:10000
----------------------	--

产品名称: MyoD (phospho Ser200) Rabbit Polyclonal Antibody
产品货号: APRab05054



蛋白分子量 (Molecular Weight) 34kDa


研究背景 (Background)

This gene encodes a nuclear protein that belongs to the basic helix-loop-helix family of transcription factors and the myogenic factors subfamily. It regulates muscle cell differentiation by inducing cell cycle arrest, a prerequisite for myogenic initiation. The protein is also involved in muscle regeneration. It activates its own transcription which may stabilize commitment to myogenesis. [provided by RefSeq, Jul 2008],function:Involved in muscle differentiation (myogenic factor). Induces fibroblasts to differentiate into myoblasts. Activates muscle-specific promoters. Interacts with and is inhibited by the twist protein. This interaction probably involves the basic domains of both proteins.,online information:MyoD entry,PTM:Acetylated by a complex containing EP300 and PCAF. The acetylation is essential to activate target genes. Conversely, its deacetylation by SIRT1 inhibits its function.,PTM:Ubiquitinated on the N-terminus; which is required for proteasomal degradation.,similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,subunit:Efficient DNA binding requires dimerization with another bHLH protein. Seems to form active heterodimers with ITF-2. Interacts with SUV39H1.,

研究领域 (Research Area)


Epigenetics and Nuclear Signaling

图片 (Image Data)



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right) , using MYOD (Phospho-Ser200) Antibody

产品名称: MyoD (phospho Ser200) Rabbit Polyclonal
Antibody
产品货号: APRab05054



Western blot analysis of lysates from Jurkat cells treated with Ca^{+} 40nM 30', using MYOD (Phospho-Ser200) Antibody.
The lane on the right is blocked with the phospho peptide.

注意事项 (Note)

For research use only.