

产品名称: NOS1 (phospho Ser852) Rabbit Polyclonal Antibody
产品货号: APRab05120



产品概述 (Summary)

产品名称 (Production Name)	NOS1 (phospho Ser852) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat,Monkey

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	NOS1
别名 (Alternative Names)	NOS1; Nitric oxide synthase; brain; Constitutive NOS; NC-NOS; NOS type I; Neuronal NOS; N-NOS; nNOS; Peptidyl-cysteine S-nitrosylase NOS1; bNOS
基因 ID (Gene ID)	4842.0
蛋白 ID (SwissProt ID)	P29475.The antiserum was produced against synthesized peptide derived from human n-NOS around the phosphorylation site of Ser852. AA range:818-867

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:10000-1:20000
----------------------	---

产品名称: NOS1 (phospho Ser852) Rabbit Polyclonal Antibody
产品货号: APRab05120

蛋白分子量 (Molecular Weight) 130-160kDa

研究背景 (Background)

The protein encoded by this gene belongs to the family of nitric oxide synthases, which synthesize nitric oxide from L-arginine. Nitric oxide is a reactive free radical, which acts as a biologic mediator in several processes, including neurotransmission, and antimicrobial and antitumoral activities. In the brain and peripheral nervous system, nitric oxide displays many properties of a neurotransmitter, and has been implicated in neurotoxicity associated with stroke and neurodegenerative diseases, neural regulation of smooth muscle, including peristalsis, and penile erection. This protein is ubiquitously expressed, with high level of expression in skeletal muscle. Multiple transcript variants that differ in the 5' UTR have been described for this gene but the full-length nature of these transcripts is not known. Additionally, alternatively spliced transcript variants encoding different isoforms/alternative products: Isoform 3 is produced by different alternative splicing events implicating either the untranslated exons TEX1 (TN-NOS) or TEX1B (TN-NOSB) leading to a N-terminus truncated protein which possesses enzymatic activity comparable to that of isoform 1. The C-terminal truncated isoform 4 is produced by insertion of the TEX2 exon between exons 3 and 4 of isoform 1, leading to a frameshift and a premature stop codon, catalytic activity: L-arginine + n NADPH + n H⁺ + m O₂ = citrulline + nitric oxide + n NADP⁺, cofactor: Binds 1 FAD, cofactor: Binds 1 FMN, cofactor: Heme group, cofactor: Tetrahydrobiopterin (BH₄). May stabilize the dimeric form of the enzyme, disease: Genetic variations in NOS1 gene are associated with susceptibility to infantile hypertrophic pyloric stenosis type 1 (IHPS1) [MIM:179010]. IHPS has an incidence of 1-5 per 1'000 live births in whites and a marked preponderance of males to females (4:1). IHPS is the most frequent disorder requiring surgery in the first year of life. The disorder is characterized by hypertrophy and hyperplasia of the circular muscle layer of the pylorus, leading to persistent vomiting 2-12 weeks after birth. Defective pyloric relaxation and increased pyloric smooth muscle mass have been suggested to be responsible for gastric-outlet obstruction, domain: The PDZ domain in the N-terminal part of the neuronal isoform participates in protein-protein interaction, and is responsible for targeting nNos to synaptic membranes in muscles, enzyme regulation: Stimulated by calcium/calmodulin. Inhibited by n-Nos-inhibiting protein (PIN) which may prevent the dimerization of the protein. Inhibited by NOSIP, function: Produces nitric oxide (NO) which is a messenger molecule with diverse functions throughout the body. In the brain and peripheral nervous system, NO displays many properties of a neurotransmitter, online information: Nitric oxide synthase entry, similarity: Belongs to the NOS family, similarity: Contains 1 FAD-binding FR-type domain, similarity: Contains 1 flavodoxin-like domain, similarity: Contains 1 PDZ (DHR) domain, subcellular location: In skeletal muscle, it is localized beneath the sarcolemma of fast-twitch muscle fiber by associating with the dystrophin glycoprotein complex. In neurons, enriched in dendritic spines, subunit: Homodimer. Interacts with DLG4; the interaction possibly being prevented by the association between NOS1 and CAPON. Forms a ternary complex with CAPON and RASD1. Forms a ternary complex with CAPON and SYN1. Interacts with ZDHHC23. Interacts with NOSIP; which may impair its synaptic location (By similarity). Interacts with HTR4, tissue specificity: Isoform 1 is ubiquitously expressed: detected in skeletal muscle and brain, also in testis, lung and kidney, and at low levels in heart, adrenal gland and retina. Not detected in the platelets. Isoform 3 is expressed only in testis. Isoform 4 is detected in testis,

产品名称: NOS1 (phospho Ser852) Rabbit Polyclonal Antibody
产品货号: AP Rab05120




skeletal muscle, lung, and kidney, at low levels in the brain, but not in the heart and adrenal gland.,


研究领域 (Research Area)

Arginine and proline metabolism; Calcium; Long-term depression; Alzheimer's disease; Amyotrophic lateral sclerosis (ALS);


图片 (Image Data)



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using n-NOS (Phospho-Ser852) Antibody




Immunofluorescence analysis of HeLa cells, using n-NOS (Phospho-Ser852) Antibody. The picture on the right is blocked with the phospho peptide.



产品名称: NOS1 (phospho Ser852) Rabbit Polyclonal Antibody
产品货号: APRab05120



Immunohistochemistry analysis of paraffin-embedded human brain, using n-NOS (Phospho-Ser852) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from A549 cells, using n-NOS (Phospho-Ser852) Antibody. The lane on the right is blocked with the phospho peptide.

注意事项 (Note)

For research use only .