

产品名称: PKC α (phospho Tyr657) Rabbit Polyclonal Antibody
产品货号: APRab05256



产品概述 (Summary)

产品名称 (Production Name)	PKC α (phospho Tyr657) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	PRKCA
别名 (Alternative Names)	PRKCA; PKCA; PRKACA; Protein kinase C alpha type; PKC-A; PKC-alpha
基因 ID (Gene ID)	5578.0
蛋白 ID (SwissProt ID)	P17252.The antiserum was produced against synthesized peptide derived from human PKC alpha around the phosphorylation site of Tyr657. AA range:623-672

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:50-1:300
蛋白分子量 (Molecular Weight)	80kDa

产品名称: PKC α (phospho Tyr657) Rabbit Polyclonal Antibody
产品货号: APRab05256




研究背景 (Background)

Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. This kinase has been reported to play roles in many different cellular processes, such as cell adhesion, cell transformation, cell cycle checkpoint, and cell volume control. Knockout studies in mice suggest that this kinase may be a fundamental regulator of cardiac contractility and Ca^{2+} handling in myocytes. [provided by RefSeq, Jul 2] catalytic activity: $\text{ATP} + \text{a protein} = \text{ADP} + \text{a phosphoprotein}$, cofactor: Binds 3 calcium ions per subunit. The ions are bound to the C2 domain, function: PKC is activated by diacylglycerol which in turn phosphorylates a range of cellular proteins. PKC also serves as the receptor for phorbol esters, a class of tumor promoters, function: This is a calcium-activated, phospholipid-dependent, serine- and threonine-specific enzyme. May play a role in cell motility by phosphorylating CSPG4, similarity: Belongs to the protein kinase superfamily, similarity: Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. PKC subfamily, similarity: Contains 1 AGC-kinase C-terminal domain, similarity: Contains 1 C2 domain, similarity: Contains 1 protein kinase domain, similarity: Contains 2 phorbol-ester/DAG-type zinc fingers, subunit: Interacts with ADAP1/CENTA1, CSPG4 and PRKCABP. Binds to SDPR in the presence of phosphatidylserine,

研究领域 (Research Area)


Regulation_Microtubule; Regulation of Actin Dynamics; Stem cell pathway; Insulin Receptor; ErbB/HER; MAPK_ERK_Growth; MAPK_G_Protein; WNT; WNT-T CELL; β -Catenin; B Cell Receptor; PI3K/Akt; mTOR; AMPK

图片 (Image Data)



Western blot analysis of lysates from COLO205 cells, using PKC alpha (Phospho-Tyr657) Antibody. The lane on the right is blocked with the phospho peptide.

产品名称: PKC α (phospho Tyr657) Rabbit Polyclonal Antibody
产品货号: APRab05256



Western Blot analysis of 293 cells using Phospho-PKC α (Y657) Polyclonal Antibody diluted at 1: 1000

注意事项 (Note)

For research use only .