

产品名称: PKC ϵ (phospho Ser729) Rabbit Polyclonal Antibody
产品货号: APRab05263



产品概述 (Summary)

产品名称 (Production Name)	PKC ϵ (phospho Ser729) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	PRKCE
别名 (Alternative Names)	PRKCE; PKCE; Protein kinase C epsilon type; nPKC-epsilon
基因 ID (Gene ID)	5581.0
蛋白 ID (SwissProt ID)	Q02156.The antiserum was produced against synthesized peptide derived from human PKC epsilon around the phosphorylation site of Ser729. AA range:688-737

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:50-1:300,ELISA 1:2000-1:20000
蛋白分子量 (Molecular Weight)	83kDa

产品名称: PKC ϵ (phospho Ser729) Rabbit Polyclonal Antibody
产品货号: APRab05263




研究背景 (Background)

protein kinase C epsilon (PRKCE) Homo sapiens Protein kinase C (PKC) is a family of serine- and threonine-specific protein kinases that can be activated by calcium and the second messenger diacylglycerol. PKC family members phosphorylate a wide variety of protein targets and are known to be involved in diverse cellular signaling pathways. PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members. This kinase has been shown to be involved in many different cellular functions, such as neuron channel activation, apoptosis, cardioprotection from ischemia, heat shock response, as well as insulin exocytosis. Knockout studies in mice suggest that this kinase is important for lipopolysaccharide (LPS)-mediated signaling in activated macrophages. activity: ATP + a protein = ADP + a phosphoprotein., domain: The C1 domain, containing the phorbol ester/DAG-type region 1 (C1A) and 2 (C1B), is the diacylglycerol sensor and the C2 domain is a non-calcium binding domain., enzyme regulation: Three specific sites; Thr-566 (activation loop of the kinase domain), Thr-710 (turn motif) and Ser-729 (hydrophobic region), need to be phosphorylated for its full activation., function: PKC is activated by diacylglycerol which in turn phosphorylates a range of cellular proteins. PKC also serves as the receptor for phorbol esters, a class of tumor promoters., function: This is calcium-independent, phospholipid-dependent, serine- and threonine-specific enzyme., PTM: Phosphorylation on Thr-566 by PDK1 triggers autophosphorylation on Ser-729., similarity: Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. PKC subfamily., similarity: Contains 1 AGC-kinase C-terminal domain., similarity: Contains 1 C2 domain., similarity: Contains 1 protein kinase domain., similarity: Contains 2 phorbol-ester/DAG-type zinc fingers., subunit: Forms a ternary complex with TRIM63 and GN2BL1.,

研究领域 (Research Area)


Regulation_Microtubule; Regulation of Actin Dynamics; Stem cell pathway; Insulin Receptor; B Cell Receptor; AMPK

图片 (Image Data)



Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using PKC epsilon (Phospho-Ser729) Antibody

产品名称: PKC ϵ (phospho Ser729) Rabbit Polyclonal Antibody
产品货号: APRab05263



Western blot analysis of lysates from HeLa cells treated with PMA 125ng/ml 30', using PKC epsilon (Phospho-Ser729) Antibody. The lane on the right is blocked with the phospho peptide.

注意事项 (Note)

For research use only.