

产品名称: PKR (phospho Thr446) Rabbit Polyclonal Antibody
产品货号: AP Rab05279



产品概述 (Summary)

产品名称 (Production Name)	PKR (phospho Thr446) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	EIF2AK2 EIF2AK2; PKR; PRKR; Interferon-induced; double-stranded RNA-activated protein kinase; Eukaryotic translation initiation factor 2-alpha kinase 2; eIF-2A protein kinase 2; Interferon-inducible RNA-dependent protein kinase; P1/eIF-2A protein k
别名 (Alternative Names)	
基因 ID (Gene ID)	5610.0
蛋白 ID (SwissProt ID)	P19525.The antiserum was produced against synthesized peptide derived from human PKR around the phosphorylation site of Thr446. AA range:413-462

产品应用 (Application)

产品名称: PKR (phospho Thr446) Rabbit Polyclonal Antibody
产品货号: APRab05279

稀释比 (Dilution Ratio) WB 1:500-1:2000, ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight) 62kDa


研究背景 (Background)

The protein encoded by this gene is a serine/threonine protein kinase that is activated by autophosphorylation after binding to dsRNA. The activated form of the encoded protein can phosphorylate translation initiation factor EIF2S1, which in turn inhibits protein synthesis. This protein is also activated by manganese ions and heparin. Three transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Oct 2011], catalytic activity: ATP + a protein = ADP + a phosphoprotein., enzyme regulation: Activity is markedly stimulated by manganese ions. Besides dsRNA, heparin is a potent activator of the kinase. Binding to dsRNA is required for dimerization leading to autophosphorylation in the activation loop and stimulation of function. Inhibited by vaccinia virus protein E3, probably via dsRNA sequestering., function: Following activation by double-stranded RNA in the presence of ATP, the kinase becomes autophosphorylated and can catalyze the phosphorylation of the translation initiation factor EIF2S1, which leads to an inhibition of the initiation of protein synthesis. Double-stranded RNA is generated during the course of a viral infection., induction: By interferon., PTM: Autophosphorylated on several Ser and Thr residues. Autophosphorylation of Thr-451 is dependent on Thr-446 and is stimulated by dsRNA binding and dimerization. Autophosphorylation apparently leads to the activation of the kinase., similarity: Belongs to the protein kinase superfamily., similarity: Belongs to the protein kinase superfamily. Ser/Thr protein kinase family. GCN2 subfamily., similarity: Contains 1 protein kinase domain., similarity: Contains 2 DRBM (double-stranded RNA-binding) domains., subunit: Homodimer. Interacts with STRBP (By similarity). Interacts with DNAJC3. Inhibited by direct interaction with viral proteins such as HCV E2, HCV NS5A and influenza A NS1. Activated by the interaction with HIV-1 Tat.,

研究领域 (Research Area)

Signal Transduction

图片 (Image Data)




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-

产品名称: PKR (phospho Thr446) Rabbit Polyclonal Antibody
产品货号: APRab05279



Phosphopeptide (Phospho-right) , using PKR (Phospho-Thr446) Antibody



Western blot analysis of lysates from NIH/3T3 cells treated with IFN 2500U/ml 30 ', using PKR (Phospho-Thr446) Antibody. The lane on the right is blocked with the phospho peptide.

注意事项 (Note)

For research use only .