

产品名称: Separase (phospho Ser801) Rabbit Polyclonal Antibody
产品货号: APRab05409



产品概述 (Summary)

产品名称 (Production Name)	Separase (phospho Ser801) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ESPL1
别名 (Alternative Names)	ESPL1; ESP1; KIAA0165; Separin; Caspase-like protein ESPL1; Extra spindle poles-like 1 protein; Separase
基因 ID (Gene ID)	9700.0
蛋白 ID (SwissProt ID)	Q14674.The antiserum was produced against synthesized peptide derived from human SEPARASE around the phosphorylation site of Ser801. AA range:767-816

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:20000-1:40000
----------------------	---

产品名称: Separase (phospho Ser801) Rabbit Polyclonal Antibody
产品货号: APRab05409



蛋白分子量 (Molecular Weight) 230kDa


研究背景 (Background)

Stable cohesion between sister chromatids before anaphase and their timely separation during anaphase are critical for chromosome inheritance. In vertebrates, sister chromatid cohesion is released in 2 steps via distinct mechanisms. The first step involves phosphorylation of STAG1 (MIM 604358) or STAG2 (MIM 300826) in the cohesin complex. The second step involves cleavage of the cohesin subunit SCC1 (RAD21; MIM 606462) by ESPL1, or separase, which initiates the final separation of sister chromatids (Sun et al., 2009 [PubMed 19345191]).[supplied by OMIM, Nov 2010],catalytic activity:All bonds known to be hydrolyzed by this endopeptidase have arginine in P1 and an acidic residue in P4. P6 is often occupied by an acidic residue or by an hydroxy-amino-acid residue, the phosphorylation of which enhances cleavage.,enzyme regulation:Regulated by at least two independent mechanisms. First, it is inactivated via its interaction with securin/PTTG1, which probably covers its active site. The association with PTTG1 is not only inhibitory, since PTTG1 is also required for activating it, the enzyme being inactive in cells in which PTTG1 is absent. PTTG1 degradation at anaphase, liberates it and triggers RAD21 cleavage. Second, phosphorylation at Ser-1126 inactivates it. The complete phosphorylation during mitosis, is removed when cells undergo anaphase. Activation of the enzyme at the metaphase-anaphase transition probably requires the removal of both securin and inhibitory phosphate.,function:Caspase-like protease, which plays a central role in the chromosome segregation by cleaving the SCC1/RAD21 subunit of the cohesin complex at the onset of anaphase. During most of the cell cycle, it is inactivated by different mechanisms.,PTM:Autocleaves. This function, which is not essential for its protease activity, is unknown.,PTM:Phosphorylated by CDC2. There are 8 Ser/Thr phosphorylation sites. Among them, Ser-1126 phosphorylation is the major site, which conducts to the enzyme inactivation.,similarity:Belongs to the peptidase C50 family.,subunit:Interacts with PTTG1. Interacts with RAD21.,

研究领域 (Research Area)

Cell_Cycle_G1S;Cell_Cycle_G2M_DNA;Oocyte meiosis;

图片 (Image Data)




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-


产品名称: Separase (phospho Ser801) Rabbit Polyclonal Antibody
产品货号: APRab05409




Phosphopeptide (Phospho-right) , using SEPARASE (Phospho-Ser801) Antibody



Immunofluorescence analysis of HUVEC cells, using SEPARASE (Phospho-Ser801) Antibody. The picture on the right is blocked with the phospho peptide.




Immunohistochemistry analysis of paraffin-embedded human colon carcinoma, using SEPARASE (Phospho-Ser801) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from 293 cells treated with EGF 200ng/ml 30 ', using SEPARASE (Phospho-Ser801) Antibody. The lane on the right is blocked with the phospho peptide.

产品名称: Separase (phospho Ser801) Rabbit Polyclonal
Antibody
产品货号: AP Rab05409



Western Blot analysis of various cells using Phospho-Separase (S801) Polyclonal Antibody diluted at 1: 1000

注意事项 (Note)

For research use only .