

产品概述 (Summary)

产品名称 (Production Name) SGK1 (phospho Ser78) Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

 应用 (Application)
 WB,IHC,ELISA

 种属反应性 (Reactivity)
 Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation) Unconjugated 修饰 (Modification) Phosphorylated

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) SGK1

SGK1; SGK; Serine/threonine-protein kinase Sgk1; Serum/glucocorticoid-别名 (Alternative Names)

regulated kinase 1

基因 ID (Gene ID) 6446.0

O00141.The antiserum was produced against synthesized peptide derived 蛋白ID (SwissProt ID)

from human SGK around the phosphorylation site of Ser78. AA range:41-90

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,IHC 1:50-1:300,ELISA 1:2000-1:20000

蛋白分子量 (Molecular Weight) 54kDa

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838



产品货号: APRab05412




研究背景 (Background)

This gene encodes a serine/threonine protein kinase that plays an important role in cellular stress response. This kinase activates certain potassium, sodium, and chloride channels, suggesting an involvement in the regulation of processes such as cell survival, neuronal excitability, and renal sodium excretion. High levels of expression of this gene may contribute to conditions such as hypertension and diabetic nephropathy. Several alternatively spliced transcript variants encoding different isoforms have been noted for this gene. [provided by RefSeq, Jan 2009], catalytic activity: ATP + a protein = ADP + a phosphoprotein.,enzyme regulation:Two specific sites, one in the kinase domain (Thr-256) and the other in the C-terminal regulatory region (Ser-422), need to be phosphorylated for its full activation. Frotein kinase that plays an important role in cellular stress response. Activates certain potassium, sodium, and chloride channels, suggesting an involvement in the regulation of processes such as cell survival, neuronal excitability, and renal sodium excretion. Sustained high levels and activity may contribute to conditions such as hypertension and diabetic nephropathy. Mediates cell survival signals, phosphorylates and negatively regulates pro-apoptotic FOXO3A. Phosphorylates NEDD4L, which leads to its inactivation and to the subsequent activation of various channels and transporters such as ENaC, Kv1.3, or EAAT1, induction: By serum and/or glucocorticoids. By excessive extracellular glucose and by TGF-beta, in cultured cells, PTM: Regulated by phosphorylation. Phosphoinositide 3-kinase (PI3-kinase) pathway promotes phosphorylation at Ser-422 which in turn increases the phosphorylation of Thr-256 by PDPK1, PTM: Ubiquitinated by NEDD4L; which promotes proteasomal degradation. Ubiquitinated by SYVN1 at the endoplasmic reticulum; which promotes rapid proteasomal degradation and maintains a high turnover rate in resting cells,,similarity:Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family, similarity: Contains 1 AGC-kinase C-terminal domain, similarity: Contains 1 protein kinase domain.,subcellular location:Nuclear, upon phosphorylation.,subunit:Interacts with NEDD4 and NEDD4L.,tissue specificity: Expressed in most tissues with highest levels in the pancreas, followed by placenta, kidney and lung.,

研究领域 (Research Area)

Aldosterone-regulated sodium reabsorption;

图片 (Image Data)




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-


E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838



Phosphopeptide (Phospho-right), using SGK (Phospho-Ser78) Antibody



Western blot analysis of lysates from 293 cells treated with UV 15 ', using SGK (Phospho-Ser78) Antibody. The lane on the right is blocked with the phospho peptide.



Western blot analysis of 293 using p-SGK1 (S78) antibody. Antibody was diluted at 1:2000

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838