

产品名称: Shc (phospho Tyr349) Rabbit Polyclonal Antibody  
产品货号: AP Rab05418



## 产品概述 (Summary)

|                        |                                                 |
|------------------------|-------------------------------------------------|
| 产品名称 (Production Name) | Shc (phospho Tyr349) Rabbit Polyclonal Antibody |
| 描述 (Description)       | Rabbit polyclonal Antibody                      |
| 宿主 (Host)              | Rabbit                                          |
| 应用 (Application)       | WB,IHC,ICC/IF,ELISA                             |
| 种属反应性 (Reactivity)     | Human,Mouse,Rat                                 |

## 产品性能 (Performance)

|                     |                                                                                                   |
|---------------------|---------------------------------------------------------------------------------------------------|
| 偶联物 (Conjugation)   | Unconjugated                                                                                      |
| 修饰 (Modification)   | Phosphorylated                                                                                    |
| 同种型 (Isotype)       | IgG                                                                                               |
| 克隆 (Clonality)      | Polyclonal                                                                                        |
| 形式 (Form)           | Liquid                                                                                            |
| 存放说明 (Storage)      | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.          |
| 储存溶液 (Buffer)       | Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N. |
| 纯化方式 (Purification) | Affinity purification                                                                             |

## 免疫原信息 (Immunogen)

|                        |                                                                                                                                                  |
|------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------|
| 基因名 (Gene Name)        | SHC1<br>SHC1; SHC; SHCA; SHC-transforming protein 1; SHC-transforming protein 3;                                                                 |
| 别名 (Alternative Names) | SHC-transforming protein A; Src homology 2 domain-containing-transforming protein C1; SH2 domain protein C1                                      |
| 基因 ID (Gene ID)        | 6464.0                                                                                                                                           |
| 蛋白 ID (SwissProt ID)   | P29353.The antiserum was produced against synthesized peptide derived from human Shc around the phosphorylation site of Tyr349. AA range:315-364 |

## 产品应用 (Application)

|                          |                                                                        |
|--------------------------|------------------------------------------------------------------------|
| 稀释比 (Dilution Ratio)     | WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000 |
| 蛋白分子量 (Molecular Weight) | 66(p66 isoform), 52(p52 isoform), 46(p46 isoform)kDa                   |

**产品名称: Shc (phospho Tyr349) Rabbit Polyclonal Antibody**  
**产品货号: AP Rab05418**



## 研究背景 (Background)


This gene encodes three main isoforms that differ in activities and subcellular location. While all three are adapter proteins in signal transduction pathways, the longest (p66Shc) may be involved in regulating life span and the effects of reactive oxygen species. The other two isoforms, p52Shc and p46Shc, link activated receptor tyrosine kinases to the Ras pathway by recruitment of the GRB2/SOS complex. p66Shc is not involved in Ras activation. Unlike the other two isoforms, p46Shc is targeted to the mitochondrial matrix. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Feb 2011], domain: In response to a variety of growth factors, isoform p46Shc and isoform p52Shc bind to phosphorylated Trk receptors through their phosphotyrosine binding (PID) and/or SH2 domains. The PID and SH2 domains bind to specific phosphorylated tyrosine residues in the Asn-Pro-Xaa-Tyr(P) motif of the Trk receptors. Isoform p46Shc and isoform p52Shc are in turn phosphorylated on three tyrosine residues within the extended proline-rich domain. These phosphotyrosines act as docking site for GRB2 and thereby are involved in Ras activation., function: Signaling adapter that couples activated growth factor receptors to signaling pathway. Isoform p46Shc and isoform p52Shc, once phosphorylated, couple activated receptor tyrosine kinases to Ras via the recruitment of the GRB2/SOS complex and are implicated in the cytoplasmic propagation of mitogenic signals. Isoform p46Shc and isoform p52Shc may thus function as initiators of the Ras signaling cascade in various non-neuronal systems. Isoform p66Shc does not mediate Ras activation, but is involved in signal transduction pathways that regulate the cellular response to oxidative stress and life span. Isoform p66Shc acts as a downstream target of the tumor suppressor p53 and is indispensable for the ability of stress-activated p53 to induce elevation of intracellular oxidants, cytochrome c release and apoptosis. The expression of isoform p66Shc has been correlated with life span., PTM: Phosphorylated by activated epidermal growth factor receptor. Isoform p46Shc and isoform p52Shc are phosphorylated on tyrosine residues of the Pro-rich domain. Isoform p66Shc is phosphorylated on Ser-36 upon treatment with insulin, hydrogen peroxide or irradiation with ultraviolet light., similarity: Contains 1 PID domain., similarity: Contains 1 SH2 domain., subcellular location: Localized to the mitochondria matrix. Targeting of isoform p46Shc to mitochondria is mediated by its first 32 amino acids, which behave as a bona fide mitochondrial targeting sequence. Isoform p52Shc and isoform p66Shc, that contain the same sequence but more internally located, display a different subcellular localization., subunit: Interacts with the Trk receptors in a phosphotyrosine-dependent manner. Interacts with the NPXY motif of tyrosine-phosphorylated IGF1R and INSR in vitro via the PID domain. Once activated, binds to GRB2. Interacts with tyrosine-phosphorylated CD3T. Interacts with the N-terminal region of APS. Interacts with phosphorylated LRP1 and IRS4. Interacts with INPP5D/SHIP1 and INPPL1/SHIP2., tissue specificity: Widely expressed. Expressed in neural stem cells but absent in mature neurons.,

## 研究领域 (Research Area)

ErbB\_HER; Chemokine; Focal adhesion; Natural killer cell mediated cytotoxicity; Neurotrophin; Insulin\_Receptor; Glioma; Chronic myeloid leukemia;


## 图片 (Image Data)

产品名称: Shc (phospho Tyr349) Rabbit Polyclonal Antibody  
产品货号: APRab05418




Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using Shc (Phospho-Tyr349) Antibody.

The picture on the right is blocked with the phospho peptide.




Western blot analysis of lysates from 293 cells treated with EGF 200ng/ml 30', using Shc (Phospho-Tyr349) Antibody.

The lane on the left is blocked with the phospho peptide.



Western Blot analysis of various cells using Phospho-Shc (Y349) Polyclonal Antibody diluted at 1: 1000

产品名称: Shc (phospho Tyr349) Rabbit Polyclonal Antibody  
产品货号: APRab05418



Immunohistochemical analysis of paraffin-embedded Human breast cancer. Antibody was diluted at 1:100 (4°,overnight) .  
High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtaned from  
antibody was pre-absorbed by immunogen peptide.

#### 注意事项 (Note)

For research use only .