

产品概述 (Summary)

产品名称 (Production Name) Sp1 (phospho Thr453) Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application)WB,IHC,ICC/IF,ELISA种属反应性 (Reactivity)Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation) Unconjugated 修饰 (Modification) Phosphorylated

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) SP1

别名 (Alternative Names) SP1; TSFP1; Transcription factor Sp1

基因 ID (Gene ID) 6667.0

P08047.The antiserum was produced against synthesized peptide derived

蛋白ID (SwissProt ID) from human SP1 around the phosphorylation site of Thr453. AA range:421-


470

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:5000-1:10000

蛋白分子量 (Molecular Weight) 90kDa

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838



产品货号: APRab05457




研究背景 (Background)

The protein encoded by this gene is a zinc finger transcription factor that binds to GC-rich motifs of many promoters. The encoded protein is involved in many cellular processes, including cell differentiation, cell growth, apoptosis, immune responses, response to DNA damage, and chromatin remodeling. Post-translational modifications such as phosphorylation, acetylation, glycosylation, and proteolytic processing significantly affect the activity of this protein, which can be an activator or a repressor. Three transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Nov 2014], function: Binds to GC box promoters elements and selectively activates mRNA synthesis from genes that contain functional recognition sites. Can interact with G/C-rich motifs from serotonin receptor promoter.,PTM:Oglycosylated; contains N-acetylglucosamine side chains, similarity; Belongs to the Sp1 C2H2-type zinc-finger protein family., similarity: Contains 3 C2H2-type zinc fingers., subunit: Interacts with ATF7IP, ATF7IP2, POGZ, HCFC1, AATF and PHC2. Interacts with varicella-zoster virus IE62 protein and HIV-1 Vpr. Interacts with SV40 VP2/3 proteins. Interacts with SV40 major capsid protein VP1; this interaction leads to a cooperativity between the two proteins in DNA binding.,


研究领域 (Research Area)

TGF-beta; Huntington's disease;

图片 (Image Data)




Immunofluorescence analysis of HeLa cells, using SP1 (Phospho-Thr453) Antibody. The picture on the right is blocked with the phospho peptide.



E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838



Immunohistochemistry analysis of paraffin-embedded human brain, using SP1 (Phospho-Thr453) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from A549 cells, using SP1 (Phospho-Thr453) Antibody. The lane on the right is blocked with the phospho peptide.

注意事项 (Note)

For research use only .

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838