

产品概述 (Summary)

产品名称 (Production Name) SSB (phospho Ser366) Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

 应用 (Application)
 WB,IHC,ICC/IF,ELISA

 种属反应性 (Reactivity)
 Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation) Unconjugated 修饰 (Modification) Phosphorylated

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) SSB

SSB; Lupus La protein; La autoantigen; La ribonucleoprotein; Sjoegren 別名 (Alternative Names)

syndrome type B antigen; SS-B

基因 ID (Gene ID) 6741.0

P05455.The antiserum was produced against synthesized peptide derived 蛋白ID (SwissProt ID)

from human SSB around the phosphorylation site of Ser366. AA range:341-390

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:20000-1:40000

蛋白分子量 (Molecular Weight) 47kDa

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838



产品货号: APRab05472




研究背景 (Background)

Sjogren syndrome antigen B(SSB) Homo sapiens The protein encoded by this gene is involved in diverse aspects of RNA metabolism, including binding and protecting poly(U) termini of nascent RNA polymerase III transcripts from exonuclease digestion, processing 5' and 3' ends of pre-tRNA precursors, acting as an RNA chaperone, and binding viral RNAs associated with hepatitis C virus. Autoantibodies reacting with this protein are found in the sera of patients with Sjogren syndrome and systemic lupus erythematosus. Alternative promoter usage results in two different transcript variants which encode the same protein. [provided by RefSeq, Jun 2014], disease: Sera from patients with systemic lupus erythematosus (SLE) often contain antibodies that react with the normal cellular La protein as if this antigen was foreign, function: La protein plays a role in the transcription of RNA polymerase III. It is most probably a transcription termination factor. Binds to the 3' termini of virtually all nascent polymerase III transcripts. It is associated with precursor forms of RNA polymerase III transcripts including tRNA and 4.5S, 5S, 7S, and 7-2 RNAs., PTM: Phosphorylated. The phosphorylation sites are at the C-terminal part of the protein., PTM: The N-terminus is blocked., similarity: Contains 1 HTH La-type RNA-binding domain., similarity: Contains 1 RRM (RNA recognition motif) domain., subunit: Interacts with DDX15. May interact with RUFY1.,

研究领域 (Research Area)

Systemic lupus erythematosus;

图片 (Image Data)




Immunofluorescence analysis of HeLa cells treated with Forskolin 40nM 30 ', using SSB (Phospho-Ser366) Antibody. The picture on the right is blocked with the phospho peptide.


Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838




产品货号: APRab05472



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using SSB (Phospho-Ser366) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from 293 cells, using SSB (Phospho-Ser366) Antibody. The lane on the right is blocked with the phospho peptide.



Immunohistochemical analysis of paraffin-embedded Human breast cancer. Antibody was diluted at 1:100 (4°, overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.

注意事项 (Note)

For research use only.