产品名称: STAM2 (phospho Tyr192) Rabbit Polyclonal

Antibody

产品货号: APRab05474



产品概述 (Summary)

产品名称 (Production Name) STAM2 (phospho Tyr192) Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application) WB,IHC,ICC/IF,ELISA

种属反应性 (Reactivity) Human, Mouse

产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Phosphorylated

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) STAM2

STAM2; HBP; Signal transducing adapter molecule 2; STAM-2; Hrs-binding 别名 (Alternative Names)

protein

基因 ID (Gene ID) 8986.0

O75886.The antiserum was produced against synthesized peptide derived

蛋白 ID (SwissProt ID) from human STAM2 around the phosphorylation site of Tyr192. AA range:161-

210

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: STAM2 (phospho Tyr192) Rabbit Polyclonal

Antibody

产品货号: APRab05474



蛋白分子量 (Molecular Weight) 58kDa


研究背景 (Background)

The protein encoded by this gene is closely related to STAM, an adaptor protein involved in the downstream signaling of cytokine receptors, both of which contain a SH3 domain and the immunoreceptor tyrosine-based activation motif (ITAM). Similar to STAM, this protein acts downstream of JAK kinases, and is phosphorylated in response to cytokine stimulation. This protein and STAM thus are thought to exhibit compensatory effects on the signaling pathway downstream of JAK kinases upon cytokine stimulation. [provided by RefSeq, Jul 2008],domain:Contains one Pro-Xaa-Val-Xaa-Leu (PxVxL) motif, which is required for interaction with chromoshadow domains. This motif requires additional residues -7, -6, +4 and +5 of the central Val which contact the chromoshadow domain.,domain:The SH3 domain mediates the interaction with USP8.,domain:The VHS and UIM domains mediate the interaction with ubiquitinated proteins.,function:Involved in intracellular signal transduction mediated by cytokines and growth factors. Upon IL-2 and GM-CSL stimulation, it plays a role in signaling leading to DNA synthesis and c-myc induction. May also play a role in T-cell development. Involved in down-regulation of receptor tyrosine kinase via multivesicular body (MVBs) when complexed with HGS (ESCRT-0 complex). The ESCRT-0 complex binds ubiquitin and acts as sorting machinery that recognizes ubiquitinated receptors and transfers them to further sequential lysosomal sorting/trafficking processes., PTM: Phosphorylated in response to IL-2, GM-CSF, EGF and PDGF, similarity: Belongs to the STAM family, similarity: Contains 1 ITAM domain, similarity: Contains 1 SH3 domain, similarity: Contains 1 UIM (ubiquitin-interacting motif) repeat, similarity: Contains 1 VHS domain., subunit: Component of the ESCRT-0 complex composed of STAM or STAM2 and HGS. Part of a complex at least composed of HSG, STAM2 and EPS15. Interacts with JAK2 and JAK3. Interacts with ubiquitinated proteins and the deubiquitinating enzyme USP8/UBPY (By similarity). Interacts (via the via the PxVxL motif) with CBX5; the interaction is direct. Interacts with VPS37C. Interacts with ubiquitin; the interaction is direct., tissue specificity: Ubiquitously expressed.,

研究领域 (Research Area)

Endocytosis;Jak_STAT;

图片 (Image Data)



Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838


产品名称: STAM2 (phospho Tyr192) Rabbit Polyclonal



产品货号: APRab05474




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using STAM2 (Phospho-Tyr192) Antibody



Western blot analysis of lysates from NIH/3T3 cells treated with EGF 200ng/ml 30 ', using STAM2 (Phospho-Tyr192)

Antibody. The lane on the right is blocked with the phospho peptide.



Western Blot analysis of 3T3 cells using Phospho-STAM2 (Y192) Polyclonal Antibody

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838