

**Antibody** 

产品货号: APRab05484



#### 产品概述 (Summary)

产品名称 (Production Name) Stat5 (phospho Ser726/731) Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application) WB,IHC,ICC/IF,ELISA 种属反应性 (Reactivity) Human, Mouse, Rat

### 产品性能 (Performance)

Unconjugated 偶联物 (Conjugation) 修饰 (Modification) Phosphorylated

同种型 (Isotype) IgG

Polyclonal 克隆 (Clonality) 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

## 免疫原信息 (Immunogen)

基因名 (Gene Name) STAT5A/STAT5B

STAT5A; STAT5; Signal transducer and activator of transcription 5A; STAT5B; 别名 (Alternative Names)

Signal transducer and activator of transcription 5B

基因 ID (Gene ID) 6777.0

P42229/P51692.The antiserum was produced against synthesized peptide

蛋白 ID (SwissProt ID) derived from human STAT5B around the phosphorylation site of Ser731. AA

range:697-746

# 产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:20000-1:40000

E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技支持) Web: https://www.enkilife.cn Tel: 027-87002838



产品货号: APRab05484



蛋白分子量 (Molecular Weight) 90kDa


## 研究背景 (Background)

The protein encoded by this gene is a member of the STAT family of transcription factors. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. This protein is activated by, and mediates the responses of many cell ligands, such as IL2, IL3, IL7 GM-CSF, erythropoietin, thrombopoietin, and different growth hormones. Activation of this protein in myeloma and lymphoma associated with a TEL/JAK2 gene fusion is independent of cell stimulus and has been shown to be essential for tumorigenesis. The mouse counterpart of this gene is found to induce the expression of BCL2L1/BCL-X(L), which suggests the antiapoptotic function of this gene in cells. Alternatively spliced transcript variants have beenfunction: Carries out a dual function: signal transduction and activation of transcription. Binds to the GAS element and activates PRL-induced transcription., online information: STAT5 entry, PTM: Tyrosine phosphorylated in response to IL-2, IL-3, IL-7, IL-15, GM-CSF, growth hormone, prolactin, erythropoietin and thrombopoietin. Tyrosine phosphorylation is required for DNA-binding activity and dimerization. Serine phosphorylation is also required for maximal transcriptional activity, similarity: Belongs to the transcription factor STAT family,,similarity:Contains 1 SH2 domain,,subcellular location:Translocated into the nucleus in response to phosphorylation, subunit: Forms a homodimer or a heterodimer with a related family member. Binds NR3C1 (By similarity). Interacts with NCOA1 and SOCS7.,

#### 研究领域(Research Area)

ErbB HER; Jak STAT; Pathways in cancer; Chronic myeloid leukemia; Acute myeloid leukemia;

## 图片 (Image Data)




Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using STAT5B (Phospho-Ser731) Antibody. The picture on the right is blocked with the phospho peptide.

E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838






产品货号: APRab05484



Western blot analysis of lysates from RAW264.7 cells treated with EGF 200ng/ml 30 ', using STAT5B (Phospho-Ser731) Antibody. The lane on the right is blocked with the phospho peptide.



Western Blot analysis of various cells using Phospho-Stat5 (S726/731) Polyclonal Antibody diluted at 1: 500

#### 注意事项 (Note)

For research use only.

E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838