产品名称: TORC1/CRTC1 (phospho-Ser151) Rabbit

Polyclonal Antibody 产品货号: APRab05571



## 产品概述 (Summary)

产品名称 (Production Name) TORC1/CRTC1 (phospho-Ser151) Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit 应用 (Application) WB

种属反应性 (Reactivity) Human, Rat, Mouse

### 产品性能 (Performance)

偶联物 (Conjugation) Unconjugated 修饰 (Modification) Phosphorylated

同种型 (Isotype) lgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

## 免疫原信息 (Immunogen)

基因名 (Gene Name) CRTC1

CREB-regulated transcription coactivator 1 (Mucoepidermoid carcinoma

别名 (Alternative Names) translocated protein 1) (Transducer of regulated cAMP response element-

binding protein 1) (TORC-1) (Transducer of CREB protein 1)

基因 ID (Gene ID) 23373.0

Q6UUV9.Synthesized phosho peptide around human TORC1 and CRTC1 蛋白ID (SwissProt ID)

(Ser151)

# 产品应用 (Application)

**稀释比 (Dilution Ratio)** WB 1:1000-1:2000

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: TORC1/CRTC1 (phospho-Ser151) Rabbit

Polyclonal Antibody 产品货号: APRab05571



蛋白分子量 (Molecular Weight) 70kDa

#### 研究背景 (Background)

disease: A chromosomal aberration involving CRTC1 is found in mucoepidermoid carcinomas, benign Warthin tumors and clear cell hidradenomas. Translocation t(11;19)(q21;p13) with MAML2. The fusion protein consists of the N-terminus of CRTC1 joined to the C-terminus of MAML2. The reciprocal fusion protein consisting of the N-terminus of MAML2 joined to the C-terminus of CRTC1 has been detected in a small number of mucoepidermoid carcinomas, function: Transcriptional coactivator for CREB1 which activates transcription through both consensus and variant cAMP response element (CRE) sites. Acts as a coactivator, in the SIK/TORC signaling pathway, being active when dephosphorylated and acts independently of CREB1 'Ser-133' phosphorylation. Enhances the interaction of CREB1 with TAF4. Regulates the expression of specific CREB-activated genes such as the steroidogenic gene, StAR. Potent coactivator of PGC1alpha and inducer of mitochondrial biogenesis in muscle cells. Also coactivator for TAX activation of the human T-cell leukemia virus type 1 (HTLV-1) long terminal repeats (LTR). In the hippocampus, involved in late-phase long-term potentiation (L-LTP) maintenance at the Schaffer collateral-CA1 synapses.,PTM:Phosphorylation/dephosphorylation states of Ser-151 are required for regulating transduction of CREB activity. TORCs are inactive when phosphorylated, and active when dephosphorylated at this site. This primary site of phosphorylation, is regulated by cAMP and calcium levels and is dependent on the phosphorylation of SIKs by LKB1 (By similarity). Phosphorylated upon DNA damage, probably by ATM or ATR., similarity: Belongs to the TORC family., subcellular location: Cytoplasmic when phosphorylated by SIK or AMPK and when sequestered by 14-3-3 proteins (By similarity). Translocated to the nucleus on Ser-151 dephosphorylation, instigated by a number of factors including calcium ion and cAMP levels., subunit: Binds, as a tetramer, through its N-terminal region, with the bZIP domain of CREB1. 'Arg-314' in the bZIP domain of CREB1 is essential for this interaction. Interaction, via its Cterminal, with TAF4, enhances recruitment of TAF4 to CREB1. Binds HTLV1 Tax., tissue specificity: Highly expressed in adult and fetal brain. Located to specific regions such as the prefrontal cortex and cerebellum. Very low expression in other tissues such as heart, spleen, lung, skeletal muscle, salivary gland, ovary and kidney.,

研究领域 (Research Area)

Signal Transduction


图片 (Image Data)

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838

产品名称: TORC1/CRTC1 (phospho-Ser151) Rabbit

Polyclonal Antibody 产品货号: APRab05571





Western Blot analysis of Hela treated or untreated by LPS lysis, using primary antibody at 1:1000 dilution. Secondary antibody was diluted at 1:10000

## 注意事项 (Note)

For research use only .

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838