

产品名称: Trk B (phospho Tyr706/Y707) Rabbit Polyclonal Antibody
产品货号: APRab05586



产品概述 (Summary)

产品名称 (Production Name)	Trk B (phospho Tyr706/Y707) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC, IF, ELISA
种属反应性 (Reactivity)	Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	NTRK2
别名 (Alternative Names)	NTRK2; TRKB; BDNF/NT-3 growth factors receptor; GP145-TrkB; Trk-B; Neurotrophic tyrosine kinase receptor type 2; TrkB tyrosine kinase; Tropomyosin-related kinase B
基因 ID (Gene ID)	4915.0
蛋白 ID (SwissProt ID)	Q16620. The antiserum was produced against synthesized peptide derived from human Trk B around the phosphorylation site of Tyr706 and Tyr707. AA range: 676-725

产品应用 (Application)

产品名称: Trk B (phospho Tyr706/Y707) Rabbit Polyclonal Antibody
产品货号: AP Rab05586

稀释比 (Dilution Ratio)

IHC 1:100-1:300, ICC/IF 1:50-1:200, ELISA 1:5000-1:20000

蛋白分子量 (Molecular Weight)


研究背景 (Background)

This gene encodes a member of the neurotrophic tyrosine receptor kinase (NTRK) family. This kinase is a membrane-bound receptor that, upon neurotrophin binding, phosphorylates itself and members of the MAPK pathway. Signalling through this kinase leads to cell differentiation. Mutations in this gene have been associated with obesity and mood disorders. Alternative splicing results in multiple transcript variants. [provided by RefSeq, May 2014], alternative products: Additional isoforms seem to exist, catalytic activity: $\text{ATP} + \text{a [protein]-L-tyrosine} = \text{ADP} + \text{a [protein]-L-tyrosine phosphate}$, function: Receptor for brain-derived neurotrophic factor (BDNF), neurotrophin-3 and neurotrophin-4/5 but not nerve growth factor (NGF). Involved in the development and/or maintenance of the nervous system. This is a tyrosine-protein kinase receptor. Known substrates for the TRK receptors are SHC1, PI-3 kinase, and PLC-gamma-1, PTM: Ligand-mediated auto-phosphorylation, similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family, similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family. Insulin receptor subfamily, similarity: Contains 1 protein kinase domain, similarity: Contains 2 Ig-like C2-type (immunoglobulin-like) domains, similarity: Contains 2 LRR (leucine-rich) repeats, subunit: Exists in a dynamic equilibrium between monomeric (low affinity) and dimeric (high affinity) structures. Binds SH2B2. Interacts with SQSTM1 and KIDINS220, tissue specificity: Isoform TrkB is widely expressed, mainly in the nervous tissue. In the CNS, expression is observed in the cerebral cortex, hippocampus, thalamus, choroid plexus, granular layer of the cerebellum, brain stem, and spinal cord. In the peripheral nervous system, it is expressed in many cranial ganglia, the ophthalmic nerve, the vestibular system, multiple facial structures, the submaxillary glands, and dorsal root ganglia. Isoform TrkB-T1 is expressed in multiple tissues, mainly in brain, pancreas, kidney and heart. Isoform TrkB-T-Shc is predominantly expressed in brain,

研究领域 (Research Area)

MAPK_ERK_Growth; MAPK_G_Protein; Neurotrophin;

图片 (Image Data)





产品名称: Trk B (phospho Tyr706/Y707) Rabbit Polyclonal
Antibody
产品货号: APRab05586

Immunohistochemistry analysis of paraffin-embedded human brain, using Trk B (Phospho-Tyr706+Tyr707) Antibody. The picture on the right is blocked with the phospho peptide.

注意事项 (Note)

For research use only .