

产品货号: APRab05595



产品概述 (Summary)

产品名称 (Production Name) Tuberin (phospho Thr1462) Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application) WB,IHC,ICC/IF,ELISA 种属反应性 (Reactivity) Human, Mouse, Rat

产品性能 (Performance)

Unconjugated 偶联物 (Conjugation) 修饰 (Modification) Phosphorylated

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) TSC2

别名 (Alternative Names) TSC2; TSC4; Tuberin; Tuberous sclerosis 2 protein

基因 ID (Gene ID) 7249.0

P49815.The antiserum was produced against synthesized peptide derived

from human Tuberin/TSC2 around the phosphorylation site of Thr1462. AA 蛋白 ID (SwissProt ID)

range:1428-1477

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:5000-1:10000

蛋白分子量 (Molecular Weight) 200kDa

E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技支持) Web: https://www.enkilife.cn Tel: 027-87002838

Antibody

产品货号: APRab05595



研究背景 (Background)

Mutations in this gene lead to tuberous sclerosis complex. Its gene product is believed to be a tumor suppressor and is able to stimulate specific GTPases. The protein associates with hamartin in a cytosolic complex, possibly acting as a chaperone for hamartin. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Jul 2008], alternative products: Additional isoforms seem to exist. Experimental confirmation may be lacking for some isoforms, disease: Defects in TSC2 are a cause of lymphangioleiomyomatosis (LAM) [MIM:606690]. LAM is a progressive and often fatal lung disease characterized by a diffuse proliferation of abnormal smooth muscle cells in the lungs. It affects almost exclusively young women and can occur as an isolated disorder or in association with tuberous sclerosis complex, disease: Defects in TSC2 are the cause of tuberous sclerosis complex (TSC) [MIM:191100]. The molecular basis of TSC is a functional impairment of the tuberin-hamartin complex. TSC is an autosomal dominant multi-system disorder that affects especially the brain, kidneys, heart, and skin. TSC is characterized by hamartomas (benign overgrowths predominantly of a cell or tissue type that occurs normally in the organ) and hamartias (developmental abnormalities of tissue combination). Clinical symptoms can range from benign hypopigmented macules of the skin to profound mental retardation with intractable seizures to premature death from a variety of disease-associated causes, function: Implicated as a tumor suppressor. May have a function in vesicular transport, but may also play a role in the regulation of cell growth arrest and in the regulation of transcription mediated by steroid receptors. Interaction between TSC1 and TSC2 may facilitate vesicular docking. Specifically stimulates the intrinsic GTPase activity of the Ras-related protein RAP1A and RAB5. Suggesting a possible mechanism for its role in regulating cellular growth. Mutations in TSC2 leads to constitutive activation of RAP1A in tumors., online information:TSC2 mutation db,PTM:Phosphorylation at Ser-1387, Ser-1418 or Ser-1420 does not affect interaction with TSC1., similarity: Contains 1 Rap-GAP domain., subcellular location: At steady state found in association with membranes, subunit: Interacts with TSC1 and HERC1; the interaction with TSC1 stabilizes TSC2 and prevents the interaction with HERC1. May also interact with the adapter molecule RABEP1. The final complex contains TSC2 and RABEP1 linked to RAB5 (Probable). Interacts with HSPA1 and HSPA8, tissue specificity: Liver, brain, heart, lymphocytes, fibroblasts, biliary epithelium, pancreas, skeletal muscle, kidney, lung and placenta.,

研究领域(Research Area)

Insulin Receptor; mTOR; B Cell Receptor; PI3K/Akt; AMPK


图片 (Image Data)

E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838




产品货号: APRab05595






Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using Tuberin/TSC2 (Phospho-Thr1462) Antibody



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma, using Tuberin/TSC2 (Phospho-Thr1462) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of 293 using p-Tuberin (T1462) antibody.

注意事项 (Note)

For research use only.

E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838