

产品名称: Myd88 (Phospho-Tyr257) Rabbit Polyclonal Antibody
产品货号: APRab05811



产品概述 (Summary)

产品名称 (Production Name)	Myd88 (Phospho-Tyr257) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB
种属反应性 (Reactivity)	Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	MYD88
别名 (Alternative Names)	Myeloid differentiation primary response protein MyD88
基因 ID (Gene ID)	4615.0
蛋白 ID (SwissProt ID)	Q99836.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000
蛋白分子量 (Molecular Weight)	33kDa

研究背景 (Background)

产品名称: Myd88 (Phospho-Tyr257) Rabbit Polyclonal Antibody
产品货号: APRab05811




disease: Defects in MYD88 are the cause of MYD88 deficiency (MYD88D) [MIM:612260]; also called recurrent pyogenic bacterial infections due to MYD88 deficiency. Patients suffer from autosomal recessive, life-threatening, often recurrent pyogenic bacterial infections, including invasive pneumococcal disease, and die between 1 and 11 months of age. Surviving patients are otherwise healthy, with normal resistance to other microbes, and their clinical status improved with age., function: Adapter protein involved in the Toll-like receptor and IL-1 receptor signaling pathway in the innate immune response. Acts via IRAK1, IRAK2 and TRAF6, leading to NF-kappa-B activation, cytokine secretion and the inflammatory response. Increases IL-8 transcription. May be involved in myeloid differentiation., similarity: Contains 1 death domain., similarity: Contains 1 TIR domain., subunit: Homodimer. Also forms heterodimers with TIRAP. Binds to TLR2, TLR4, IRAK1 and IRAK2 via their respective TIR domains. Interacts with IL1RL1., tissue specificity: Ubiquitous.,

研究领域 (Research Area)

Signal Transduction

图片 (Image Data)



Western Blot analysis of 1 A431 cell, 2 Serum-free treated, using primary antibody at 1:1000 dilution. Secondary antibody was diluted at 1:10000

注意事项 (Note)

For research use only.