

产品名称: Smad3 (Phospho-Ser213) Rabbit Polyclonal Antibody
产品货号: APRab06081



产品概述 (Summary)

产品名称 (Production Name)	Smad3 (Phospho-Ser213) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Phosphorylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	SMAD3 MADH3
别名 (Alternative Names)	Mothers against decapentaplegic homolog 3 (MAD homolog 3) (Mad3) (Mothers against DPP homolog 3) (hMAD-3) (JV15-2) (SMAD family member 3) (SMAD 3) (Smad3) (hSMAD3)
基因 ID (Gene ID)	4088.0
蛋白 ID (SwissProt ID)	P84022.Synthesized phospho derived from human Smad3 (Phospho-Ser213)

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:50-1:300,ELISA 1:2000-1:20000
蛋白分子量 (Molecular Weight)	50kDa

产品名称: Smad3 (Phospho-Ser213) Rabbit Polyclonal Antibody
产品货号: APRab06081




研究背景 (Background)

The protein encoded by this gene belongs to the SMAD, a family of proteins similar to the gene products of the *Drosophila* gene *mothers against decapentaplegic* (Mad) and the *C. elegans* gene *Sma*. SMAD proteins are signal transducers and transcriptional modulators that mediate multiple signaling pathways. This protein functions as a transcriptional modulator activated by transforming growth factor-beta and is thought to play a role in the regulation of carcinogenesis. [provided by RefSeq, Apr 2009], disease: Defects in SMAD3 may be a cause of colorectal cancer (CRC) [MIM:114500], domain: The MH2 domain is sufficient to carry protein nuclear export, function: Transcriptional modulator activated by TGF-beta (transforming growth factor) and activin type 1 receptor kinase. SMAD3 is a receptor-regulated SMAD (R-SMAD), PTM: Phosphorylated on serine by TGF-beta and activin type 1 receptor kinases, similarity: Belongs to the dwarfin/SMAD family, similarity: Contains 1 MH1 (MAD homology 1) domain, similarity: Contains 1 MH2 (MAD homology 2) domain, subcellular location: In the cytoplasm in the absence of ligand. Migration to the nucleus when complexed with Smad4, subunit: Interacts with HGS. Interacts with NEDD4L in response to TGF-beta. Interacts with TTRAP (By similarity). Interacts with SARA (SMAD anchor for receptor activation); form trimers with another SMAD3 and the co-SMAD SMAD4. Interacts with JUN/FOS, vitamin D receptor, homeobox protein TGIF and TGIF2, PEBP2-alpha C subunit, CREB-binding protein (CBP), p300, SKI, SNON, ATF2, SMURF2, AIP1, DACH1 and TGFB11. Part of a complex consisting of AIP1, ACVR2A, ACVR1B and SMAD3. Found in a complex with SMAD2 and TRIM33 upon addition of TGF-beta. Interacts with SMAD2 and TRIM33. Found in a complex with SMAD3, Ran and XPO4. Interacts with XPO4. Interacts with LBXCOR1 and CORL2,

研究领域 (Research Area)

Cell_Cycle_G1S; Cell_Cycle_G2M_DNA; WNT; WNT-T CELL; TGF-beta; Adherens_Junction; Pathways in cancer; Colorectal cancer; Pancreatic cancer; Chronic myeloid leukemia;

图片 (Image Data)



Western blot analysis of various lysate, antibody was diluted at 1000. Secondary antibody was diluted at 1:20000

注意事项 (Note)

**产品名称: Smad3 (Phospho-Ser213) Rabbit Polyclonal
Antibody**
产品货号: APRab06081



For research use only .