

产品货号: APRab06180

产品概述 (Summary)

产品名称 (Production Name) Cytokeratin 8 (Acetyl Lys483) Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application)WB,IHC,ICC/IF,ELISA种属反应性 (Reactivity)Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation) Unconjugated **修饰 (Modification**) Acetylated

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) KRT8

KRT8; CYK8; Keratin, type II cytoskeletal 8; Cytokeratin-8; CK-8; Keratin-8; K8; **别名 (Alternative Names)**

Type-II keratin Kb8

基因 ID (Gene ID) 3856.0

P05787.The antiserum was produced against synthesized Acetyl-peptide

蛋白ID (SwissProt ID) derived from human K8 around the Acetylation site of Lys483. AA range:434-

483

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: Cytokeratin 8 (Acetyl Lys483) Rabbit Polyclonal

Antibody

产品货号: APRab06180



蛋白分子量 (Molecular Weight) 53kDa


研究背景 (Background)

keratin 8(KRT8) Homo sapiens This gene is a member of the type II keratin family clustered on the long arm of chromosome 12. Type I and type II keratins heteropolymerize to form intermediate-sized filaments in the cytoplasm of epithelial cells. The product of this gene typically dimerizes with keratin 18 to form an intermediate filament in simple single-layered epithelial cells. This protein plays a role in maintaining cellular structural integrity and also functions in signal transduction and cellular differentiation. Mutations in this gene cause cryptogenic cirrhosis. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jan 2012], disease: Defects in KRT8 are a cause of cryptogenic cirrhosis [MIM:215600], function: Together with KRT19, helps to link the contractile apparatus to dystrophin at the costameres of striated muscle., miscellaneous: There are two types of cytoskeletal and microfibrillar keratin: I (acidic; 40-55 kDa) and II (neutral to basic; 56-70 kDa), PTM: O-glycosylated at multiple sites; glycans consist of single Nacetylglucosamine residues., PTM: Phosphorylation on serine residues is enhanced during EGF stimulation and mitosis. Ser-74 phosphorylation plays an important role in keratin filament reorganization., similarity: Belongs to the intermediate filament family, subunit: Heterotetramer of two type I and two type II keratins. keratin-8 associates with keratin-18. Associates with KRT20. Interacts with HCV core protein and PNN. When associated with KRT19, interacts with DMD. Interacts with TCHP, tissue specificity: Observed in muscle fibers accumulating in the costameres of myoplasm at the sarcolemma membrane in structures that contain dystrophin and spectrin. Expressed in gingival mucosa and hard palate of the oral cavity.,

研究领域(Research Area)

Signal Transduction

图片 (Image Data)



Western blot analysis of lysate from A549 cells, using K8 (Acetyl-Lys483) Antibody.


E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838




Antibody

产品货号: APRab06180






Western Blot analysis of A549 cells using Acetyl-Cytokeratin 8 (K483) Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Western Blot analysis of A549 cells using Acetyl-Cytokeratin 8 (K483) Polyclonal Antibody. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-breast, antibody was diluted at 1:100

注意事项 (Note)

For research use only.

E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838