

产品名称: Histone H4 (Acetyl Lys12) Rabbit Polyclonal Antibody
产品货号: APRab06212



产品概述 (Summary)

产品名称 (Production Name)	Histone H4 (Acetyl Lys12) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat,Monkey

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Acetylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	HIST1H4A
别名 (Alternative Names)	HIST1H4A; H4/A; H4FA; HIST1H4B; H4/I; H4FI; HIST1H4C; H4/G; H4FG; HIST1H4D; H4/B; H4FB; HIST1H4E; H4/J; H4FJ; HIST1H4F; H4/C; H4FC; HIST1H4H; H4/H; H4FH; HIST1H4I; H4/M; H4FM; HIST1H4J; H4/E; H4FE; HIST1H4K; H4/D; H4FD; HIST1H4L; H4/K; H4FK;H4K12AC
基因 ID (Gene ID)	121504/554313/8294/8359/8360/8361/8362/8363/8364/8365/8366/8367/8368/8370
蛋白 ID (SwissProt ID)	P62805.The antiserum was produced against synthesized peptide derived from human Histone H4 around the acetylated site of Lys12. AA range:10-59

产品应用 (Application)

产品名称: Histone H4 (Acetyl Lys12) Rabbit Polyclonal Antibody
产品货号: APRab06212



稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	11kDa

研究背景 (Background)


Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form an octamer, around which approximately 146 bp of DNA is wrapped in repeating units, called nucleosomes. The linker histone, H1, interacts with linker DNA between nucleosomes and functions in the compaction of chromatin into higher order structures. This gene is intronless and encodes a replication-dependent histone that is a member of the histone H4 family. Transcripts from this gene lack polyA tails but instead contain a palindromic termination element. This gene is found in the histone microcluster on chromosome 6p21.33. [provided by RefSeq, Aug 2015],function:Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.,PTM:Acetylation at Lys-6, Lys-9, Lys-13 and Lys-17 occurs in coding regions of the genome but not in heterochromatin.,PTM:Citrullination at Arg-4 by PADI4 impairs methylation.,PTM:Monomethylated, dimethylated or trimethylated at Lys-21. Monomethylation is performed by SET8. Trimethylation is performed by SUV420H1 and SUV420H2 and induces gene silencing.,PTM:Monomethylation at Arg-4 by PRMT1 favors acetylation at Lys-9 and Lys-13. Demethylation is performed by JMJD6.,PTM:Sumoylated, which is associated with transcriptional repression.,PTM:Ubiquitinated by the CUL4-DDB-RBX1 complex in response to ultraviolet irradiation. This may weaken the interaction between histones and DNA and facilitate DNA accessibility to repair proteins.,similarity:Belongs to the histone H4 family.,subunit:The nucleosome is a histone octamer containing two molecules each of H2A, H2B, H3 and H4 assembled in one H3-H4 heterotetramer and two H2A-H2B heterodimers. The octamer wraps approximately 147 bp of DNA.,

研究领域 (Research Area)


Protein_Acetylation

图片 (Image Data)


产品名称: Histone H4 (Acetyl Lys12) Rabbit Polyclonal Antibody
产品货号: APRab06212



Immunofluorescence analysis of HeLa cells, using Histone H4 (Acetyl-Lys12) Antibody. The picture on the right is blocked with the synthesized peptide.




Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using Histone H4 (Acetyl-Lys12) Antibody. The picture on the right is blocked with the synthesized peptide.




Western blot analysis of lysates from COS7 cells, treated with TSA 400nM 24h, using Histone H4 (Acetyl-Lys12) Antibody. The lane on the right is blocked with the synthesized peptide.


产品名称: Histone H4 (Acetyl Lys12) Rabbit Polyclonal Antibody
产品货号: AP Rab06212



Western Blot analysis of KB cells using Acetyl-Histone H4 (K12) Polyclonal Antibody diluted at 1: 500. Secondary antibody was diluted at 1:20000



Western Blot analysis of 293 cells using Acetyl-Histone H4 (K12) Polyclonal Antibody diluted at 1: 500. Secondary antibody was diluted at 1:20000



Western blot analysis of Mouse-kidney HELA lysis using Acetyl-Histone H4 (K12) antibody. Antibody was diluted at 1:500.
Secondary antibody was diluted at 1:20000

注意事项 (Note)

For research use only .