

产品名称: HMG-1 (Acetyl Lys12) Rabbit Polyclonal Antibody
产品货号: APRab06217



产品概述 (Summary)

产品名称 (Production Name)	HMG-1 (Acetyl Lys12) Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Acetylated
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	HMGB1
别名 (Alternative Names)	HMGB1; HMG1; High mobility group protein B1; High mobility group protein 1; HMG-1
基因 ID (Gene ID)	3146.0
蛋白 ID (SwissProt ID)	P09429.Synthesized acetyl-peptide derived from the N-terminal region of human HMG-1 around the acetylation site of K12.

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	about 30kDa

产品名称: HMG-1 (Acetyl Lys12) Rabbit Polyclonal Antibody
产品货号: APRab06217


研究背景 (Background)

This gene encodes a protein that belongs to the High Mobility Group-box superfamily. The encoded non-histone, nuclear DNA-binding protein regulates transcription, and is involved in organization of DNA. This protein plays a role in several cellular processes, including inflammation, cell differentiation and tumor cell migration. Multiple pseudogenes of this gene have been identified. Alternative splicing results in multiple transcript variants that encode the same protein. [provided by RefSeq, Sep 2015],function: Binds preferentially single-stranded DNA and unwinds double stranded DNA.,similarity: Belongs to the HMGB family.,similarity: Contains 2 HMG box DNA-binding domains.,


研究领域 (Research Area)

Base excision repair;

图片 (Image Data)




Western Blot analysis of HepG2 cells treated with UV using Acetyl-HMG-1 (K12) Polyclonal Antibody. Antibody was diluted at 1:1000. Secondary antibody was diluted at 1:20000




Immunohistochemical analysis of paraffin-embedded human breast, antibody was diluted at 1:100

产品名称: HMG-1 (Acetyl Lys12) Rabbit Polyclonal Antibody
产品货号: APRab06217



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:100

注意事项 (Note)

For research use only .