

产品概述 (Summary)

产品名称 (Production Name) p73 (Acetyl Lys321) Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application)WB,IHC,ICC/IF,ELISA种属反应性 (Reactivity)Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation) Unconjugated 修饰 (Modification) Acetylated

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) TP73

TP73; P73; Tumor protein p73; p53-like transcription factor; p53-related **别名 (Alternative Names)**

protein

基因 ID (Gene ID) 7161.0

O15350.The antiserum was produced against synthesized Acetyl-peptide

蛋白ID (SwissProt ID) derived from human p73 around the Acetylation site of Lys321. AA range:281-

330

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000

蛋白分子量 (Molecular Weight) 70kDa

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: p73 (Acetyl Lys321) Rabbit Polyclonal Antibody EnkiLife 产品货号: APRab06247

研究背景 (Background)

tumor protein p73(TP73) Homo sapiens This gene encodes a member of the p53 family of transcription factors involved in cellular responses to stress and development. It maps to a region on chromosome 1p36 that is frequently deleted in neuroblastoma and other tumors, and thought to contain multiple tumor suppressor genes. The demonstration that this gene is monoallelically expressed (likely from the maternal allele), supports the notion that it is a candidate gene for neuroblastoma. Many transcript variants resulting from alternative splicing and/or use of alternate promoters have been found for this gene, but the biological validity and the full-length nature of some variants have not been determined. [provided by RefSeq, Feb 2011], cofactor: Binds 1 zinc ion per subunit, disease: Maps to a chromosome region frequently mutated in diverse cell lines of human cancer. Appears not to be frequently mutated in human cancers, in contrast to p53. Hemizygosity is observed in neuroblastoma and oligodendroglioma.,domain:Possesses an acidic transactivation domain, a central DNA binding domain and a C-terminal oligomerization domain that binds to the ABL tyrosine kinase SH3 domain, domain: The WW-binding motif mediates interaction with WWOX, function: Participates in the apoptotic response to DNA damage. Isoforms containing the transactivation domain are pro-apoptotic, isoforms lacking the domain are antiapoptotic and block the function of p53 and transactivating p73 isoforms. May be a tumor suppressor protein, induction: Not induced by DNA damage. Isoforms lacking the transactivation domain block gene induction, miscellaneous: Activated and stabilized by interaction with RANBP9, PTM: Isoform alpha (but not isoform beta) is sumoylated on Lys-627, which potentiates proteasomal degradation but does not affect transcriptional activity, similarity: Belongs to the p53 family, similarity: Contains 1 SAM (sterile alpha motif) domain, subcellular location: Accumulates in the nucleus in response to DNA damage, subunit: Found in a complex with p53/TP53 and CABLES1. The C-terminal oligomerization domain binds to the ABL tyrosine kinase SH3 domain. Interacts with HECW2. Isoform Beta interacts homotypically and with p53/TP53, whereas isoform Alpha does not. Isoform Gamma interacts homotypically and with all p73 isoforms. Isoform Delta interacts with isoform Gamma, isoform Alpha, and homotypically. Isoforms Alpha and Beta interact with HIPK2. Isoform Alpha interacts with RANBP9. Isoform Beta interacts with WWOX.,tissue specificity:Brain, kidney, placenta, colon, heart, liver, spleen, skeletal muscle, prostate, thymus and pancreas. Highly expressed in fetal tissue.,


研究领域 (Research Area)

p53;Neurotrophin;

图片 (Image Data)

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838






Western blot analysis of lysate from K562 cells, using p73 (Acetyl-Lys321) Antibody.



Western Blot analysis of K562 cells using Acetyl-p73 (K321) Polyclonal Antibody.. Secondary antibody was diluted at 1:20000




Western Blot analysis of K562 cells using Acetyl-p73 (K321) Polyclonal Antibody. Secondary antibody was diluted at 1:20000






产品货号: APRab06247




Immunohistochemical analysis of paraffin-embedded human-breast, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-breast, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:100



E-mail: order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Web: https://www.enkilife.cn Tel: 027-87002838



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:100

注意事项 (Note)

For research use only .

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838