

产品名称: 14-3-3 ζ/δ Rabbit Polyclonal Antibody
产品货号: APRab06283



产品概述 (Summary)

产品名称 (Production Name)	14-3-3 ζ/δ Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	YWHAZ
别名 (Alternative Names)	YWHAZ; 14-3-3 protein zeta/delta; Protein kinase C inhibitor protein 1; KCIP-1
基因 ID (Gene ID)	7534.0
蛋白 ID (SwissProt ID)	P63104.The antiserum was produced against synthesized peptide derived from human 14-3-3 zeta/delta. AA range:196-245

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	28kDa

研究背景 (Background)

产品名称: 14-3-3 ζ/δ Rabbit Polyclonal Antibody
产品货号: APRab06283




This gene product belongs to the 14-3-3 family of proteins which mediate signal transduction by binding to phosphoserine-containing proteins. This highly conserved protein family is found in both plants and mammals, and this protein is 99% identical to the mouse, rat and sheep orthologs. The encoded protein interacts with IRS1 protein, suggesting a role in regulating insulin sensitivity. Several transcript variants that differ in the 5' UTR but that encode the same protein have been identified for this gene. [provided by RefSeq, Oct 2008], caution: Was originally (PubMed:1577711) thought to have phospholipase A2 activity, function: Adapter protein implicated in the regulation of a large spectrum of both general and specialized signaling pathway. Binds to a large number of partners, usually by recognition of a phosphoserine or phosphothreonine motif. Binding generally results in the modulation of the activity of the binding partner, PTM: The delta, brain-specific form differs from the zeta form in being phosphorylated (By similarity). Phosphorylation on Ser-184 by MAPK8; promotes dissociation of BAX and translocation of BAX to mitochondria. Phosphorylation on Ser-58 by PKA; disrupts homodimerization and heterodimerization with YHAE and TP53. This phosphorylation appears to be activated by sphingosine. Phosphorylation on Thr-232; inhibits binding of RAF1, similarity: Belongs to the 14-3-3 family, subcellular location: Located to stage I to stage IV melanosomes, subunit: Homodimer. Heterodimerizes with YWHAE. Homo- and hetero-dimerization is inhibited by phosphorylation on Ser-58. Interacts with FOXO4, NOXA1, SSH1 and ARHGEF2. Interacts with PCK1 and BSPRY (By similarity). Interacts with WEE1 (C-terminal) (By similarity). Interacts with MLF1 (phosphorylated form); the interaction retains it in the cytoplasm (By similarity). Interacts with Thr-phosphorylated ITGB2 (By similarity). Interacts with Pseudomonas aeruginosa exoS (unphosphorylated form). Interacts with BAX; the interaction occurs in the cytoplasm. Under stress conditions, MAPK8-mediated phosphorylation releases BAX to mitochondria. Interacts with phosphorylated RAF1; the interaction is inhibited when YWHAZ is phosphorylated on Thr-232. Interacts with TP53; the interaction enhances p53 transcriptional activity. The Ser-58 phosphorylated form inhibits this interaction and p53 transcriptional activity. Interacts with ABL1 (phosphorylated form); the interaction retains ABL1 in the cytoplasm. Interacts with AANAT ('Thr-31' phosphorylated form); the interaction modulates AANAT enzymatic activity through preventing dephosphorylation and/or proteolysis and stabilizing substrate binding. Subsequently, a second molecule of AANAT ('Ser-205' phosphorylated form), can bind the other YWHAZ monomer with similar effect. Interacts with AKT1; the interaction phosphorylates YWHAZ and modulates dimerization.

研究领域 (Research Area)


Cell_Cycle_G1S; Cell_Cycle_G2M_DNA; Oocyte meiosis; Neurotrophin; Pathogenic Escherichia coli infection;

图片 (Image Data)


产品名称: 14-3-3 ζ/δ Rabbit Polyclonal Antibody
产品货号: APRab06283



Immunofluorescence analysis of NIH/3T3 cells, using 14-3-3 zeta/delta Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using 14-3-3 zeta/delta Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from K562 cells, using 14-3-3 zeta/delta Antibody. The lane on the right is blocked with the synthesized peptide.

注意事项 (Note)

For research use only .