

产品名称: ACAD-9 Rabbit Polyclonal Antibody
产品货号: APRab06460



产品概述 (Summary)

产品名称 (Production Name)	ACAD-9 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC, IF, ELISA
种属反应性 (Reactivity)	Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ACAD9
别名 (Alternative Names)	ACAD9; Acyl-CoA dehydrogenase family member 9; mitochondrial; ACAD-9
基因 ID (Gene ID)	28976.0
蛋白 ID (SwissProt ID)	Q9H845.Synthesized peptide derived from ACAD-9 . at AA range: 530-610

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:100-1:300, ICC/IF 1:50-1:200, ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	

研究背景 (Background)

This gene encodes a member of the acyl-CoA dehydrogenase family. Members of this family of proteins localize to the


产品名称: ACAD-9 Rabbit Polyclonal Antibody
产品货号: APRab06460



mitochondria and catalyze the rate-limiting step in the beta-oxidation of fatty acyl-CoA. The encoded protein is specifically active toward palmitoyl-CoA and long-chain unsaturated substrates. Mutations in this gene cause acyl-CoA dehydrogenase family member type 9 deficiency. Alternate splicing results in multiple transcript variants.[provided by RefSeq, Mar 2010],cofactor:FAD.,disease:Defects in ACAD9 are a cause of acyl-CoA dehydrogenase family member type 9 deficiency (ACAD9 deficiency) [MIM:611126]. ACAD9 deficiency patients present with episodic liver dysfunction during otherwise mild illnesses or cardiomyopathy, along with chronic neurologic dysfunction.,function:Has a dehydrogenase activity on palmitoyl-CoA (C16:0) and stearoyl-CoA (C18:0). It is three times more active on palmitoyl-CoA than on stearoyl-CoA. Has little activity on octanoyl-CoA (C8:0), butyryl-CoA (C4:0) or isovaleryl-CoA (5:0).,similarity:Belongs to the acyl-CoA dehydrogenase family.,tissue specificity:Ubiquitously expressed in most normal human tissues and cancer cell lines with high level of expression in heart, skeletal muscles, brain, kidney and liver.,

研究领域 (Research Area)

图片 (Image Data)



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200 (4° overnight) . 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 30min) .

注意事项 (Note)

For research use only .