产品名称: ACBP Rabbit Polyclonal Antibody

产品货号: APRab06477



产品概述 (Summary)

产品名称 (Production Name) ACBP Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit 应用 (Application) WB,ELISA

种属反应性 (Reactivity) Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) DBI

DBI; Acyl-CoA-binding protein; ACBP; Diazepam-binding inhibitor; DBI; **别名 (Alternative Names)**

Endozepine; EP

基因 ID (Gene ID) 1622.0

P07108.The antiserum was produced against synthesized peptide derived 蛋白ID (SwissProt ID)

from the C-terminal region of human DBI. AA range:38-87

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,ELISA 1:10000-1:20000

蛋白分子量 (Molecular Weight) 10kDa

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: ACBP Rabbit Polyclonal Antibody

产品货号: APRab06477




研究背景 (Background)

This gene encodes diazepam binding inhibitor, a protein that is regulated by hormones and is involved in lipid metabolism and the displacement of beta-carbolines and benzodiazepines, which modulate signal transduction at type A gamma-aminobutyric acid receptors located in brain synapses. The protein is conserved from yeast to mammals, with the most highly conserved domain consisting of seven contiguous residues that constitute the hydrophobic binding site for medium- and long-chain acyl-Coenzyme A esters. Diazepam binding inhibitor is also known to mediate the feedback regulation of pancreatic secretion and the postprandial release of cholecystokinin, in addition to its role as a mediator in corticotropin-dependent adrenal steroidogenesis. Three pseudogenes located on chromosomes 6, 8 and 16 have been identified. Multiple transcript variants encoding different isoformfunction:Binds medium- and long-chain acyl-CoA esters with very high affinity and may function as an intracellular carrier of acyl-CoA esters. It is also able to displace diazepam from the benzodiazepine (BZD) recognition site located on the GABA type A receptor. It is therefore possible that this protein also acts as a neuropeptide to modulate the action of the GABA receptor.,similarity:Belongs to the ACBP family,,similarity;Contains 1 ACB (acyl-CoA-binding) domain,,subunit:Monomer.,


研究领域 (Research Area)

PPAR;

图片 (Image Data)



Western blot analysis of lysate from MCF7 cells, using DBI Antibody.



Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838



产品货号: APRab06477



Western Blot analysis of MCF7, U251, mouse liver, mouse kidney cells using ACBP Polyclonal Antibody.. Secondary antibody was diluted at 1:20000

注意事项 (Note)

For research use only .

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838