

产品名称: ACK Rabbit Polyclonal Antibody
产品货号: APRab06504



产品概述 (Summary)

产品名称 (Production Name)	ACK Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB, ICC/IF, ELISA
种属反应性 (Reactivity)	Human, Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	TNK2
别名 (Alternative Names)	TNK2; ACK1; Activated CDC42 kinase 1; ACK-1; Tyrosine kinase non-receptor protein 2
基因 ID (Gene ID)	10188.0
蛋白 ID (SwissProt ID)	Q07912. The antiserum was produced against synthesized peptide derived from human ACK1. AA range: 250-299

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000, ICC/IF 1:50-1:200, ELISA 1:5000-1:20000
蛋白分子量 (Molecular Weight)	115kDa

产品名称: ACK Rabbit Polyclonal Antibody
产品货号: APRab06504




研究背景 (Background)

This gene encodes a tyrosine kinase that binds Cdc42Hs in its GTP-bound form and inhibits both the intrinsic and GTPase-activating protein (GAP)-stimulated GTPase activity of Cdc42Hs. This binding is mediated by a unique sequence of 47 amino acids C-terminal to an SH3 domain. The protein may be involved in a regulatory mechanism that sustains the GTP-bound active form of Cdc42Hs and which is directly linked to a tyrosine phosphorylation signal transduction pathway. Several alternatively spliced transcript variants have been identified from this gene, but the full-length nature of only two transcript variants has been determined. [provided by RefSeq, Jul 2008], catalytic activity: ATP + a [protein]-L-tyrosine = ADP + a [protein]-L-tyrosine phosphate., cofactor: Magnesium., enzyme regulation: The SH3 domain appears to play an autoinhibitory role., function: Downstream effector of CDC42 which mediates CDC42-dependent cell migration via phosphorylation of BCAR1. Binds to both poly- and mono-ubiquitin and regulates ligand-induced degradation of EGFR. Participates in clathrin-mediated endocytosis. May be involved both in adult synaptic function and plasticity and in brain development., sequence caution: Unlikely isoform. Aberrant splice sites., similarity: Belongs to the protein kinase superfamily. Tyr protein kinase family., similarity: Contains 1 CRIB domain., similarity: Contains 1 protein kinase domain., similarity: Contains 1 SH3 domain., subunit: Interacts with CDC42. Interacts with activated CSPG4.,


研究领域 (Research Area)

图片 (Image Data)



Western blot analysis of the lysates from COLO205 cells using ACK1 antibody.

产品名称: ACK Rabbit Polyclonal Antibody
产品货号: APRab06504



Western Blot analysis of A549 cells using ACK Polyclonal Antibody. Secondary antibody was diluted at 1:20000

注意事项 (Note)

For research use only .