

产品名称: ADAMTS-1 Rabbit Polyclonal Antibody
产品货号: APRab06595



产品概述 (Summary)

产品名称 (Production Name)	ADAMTS-1 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ADAMTS1
别名 (Alternative Names)	ADAMTS1; KIAA1346; METH1; A disintegrin and metalloproteinase with thrombospondin motifs 1; ADAM-TS 1; ADAM-TS1; ADAMTS-1; METH-1
基因 ID (Gene ID)	9510.0
蛋白 ID (SwissProt ID)	Q9UHI8.Synthesized peptide derived from ADAMTS-1 . at AA range: 160-240

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:5000-1:10000
蛋白分子量 (Molecular Weight)	105kDa

研究背景 (Background)


产品名称: ADAMTS-1 Rabbit Polyclonal Antibody
产品货号: APRab06595



This gene encodes a member of the ADAMTS (a disintegrin and metalloproteinase with thrombospondin motif) protein family. Members of the family share several distinct protein modules, including a propeptide region, a metalloproteinase domain, a disintegrin-like domain, and a thrombospondin type 1 (TS) motif. Individual members of this family differ in the number of C-terminal TS motifs, and some have unique C-terminal domains. The protein encoded by this gene contains two disintegrin loops and three C-terminal TS motifs and has anti-angiogenic activity. The expression of this gene may be associated with various inflammatory processes as well as development of cancer cachexia. This gene is likely to be necessary for normal growth, fertility, and organ morphology and function. [provided by RefSeq, Jul 2008], catalytic activity: Cleaves aggrecan at the 1938-Glu-|-Leu-1939 site, within the chondroitin sulfate attachment domain., cofactor: Binds 1 zinc ion per subunit., domain: The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme., domain: The spacer domain and the TSP type-1 domains are important for a tight interaction with the extracellular matrix., function: Cleaves aggrecan, a cartilage proteoglycan, and may be involved in its turnover (By similarity). Has angiogenic inhibitor activity. Active metalloprotease, which may be associated with various inflammatory processes as well as development of cancer cachexia. May play a critical role in follicular rupture., PTM: The precursor is cleaved by a furin endopeptidase., similarity: Contains 1 disintegrin domain., similarity: Contains 1 peptidase M12B domain., similarity: Contains 3 TSP type-1 domains.,


研究领域 (Research Area)

图片 (Image Data)



Western Blot analysis of various cells using ADAMTS-1 Polyclonal Antibody

产品名称: ADAMTS-1 Rabbit Polyclonal Antibody
产品货号: APRab06595



Western Blot analysis of HUVEC cells using ADAMTS-1 Polyclonal Antibody

注意事项 (Note)

For research use only .