

产品名称: AID Rabbit Polyclonal Antibody

产品货号: APRab06697

产品概述 (Summary)

描述(Description)	Rabbit polyclonal Antibody
宿主(Host)	Rabbit
应用(Application)	WB,IHC,ICC/IF,ELISA
种属反应(Reactivity)	Human,Mouse,Rat
偶联物(Conjugation)	Unconjugated
修饰(Modification)	Unmodified
同种型(Isotype)	IgG
克隆(Clonality)	Polyclonal
剂型(Form)	Liquid
保存条件(Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液(Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化 (Purification)	Affinity purification

产品应用(Application)

稀释比(Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:20000
分子量(Molecular Weight)	24kDa

抗原信息(Antigen Information)

基因名(Gene Name)	AICDA
别名(Alternative Names)	AICDA; AID; Activation-induced cytidine deaminase; Cytidine aminohydrolase
基因 ID(Gene ID)	57379.0
SwissProt ID	Q9GZX7
免疫原(Immunogen)	The antiserum was produced against synthesized peptide derived from the Internal region of human AICDA. AA range:81-130

研究背景 (Background)

This gene encodes a RNA-editing deaminase that is a member of the cytidine deaminase family. The protein is involved in somatic hypermutation, gene conversion, and class-switch recombination of immunoglobulin genes. Defects in this gene are

the cause of autosomal recessive hyper-IgM immunodeficiency syndrome type 2 (HIGM2). [provided by RefSeq, Feb 2009],catalytic activity:Cytidine + H(2)O = uridine + NH(3),cofactor:Zinc,disease:Defects in AICDA are the cause of autosomal recessive hyper-IgM immunodeficiency syndrome type 2 (HIGM2) [MIM:605258]. HIGM2 is characterized by normal or elevated serum IgM levels with absence of IgG, IgA, and IgE, resulting in a profound susceptibility to bacterial infections. HIGM2 causes the absence of Ig class switch recombination (CSR), the lack of Ig somatic hypermutations, and lymph node hyperplasia caused by the presence of giant germinal centers.,function:RNA-editing deaminase involved in somatic hypermutation, gene conversion, and class-switch recombination. Required for several crucial steps of B-cell terminal differentiation necessary for efficient antibody responses.,online information:AICDA mutation db,similarity:Belongs to the cytidine and deoxycytidylate deaminase family.,tissue specificity:Strongly expressed in lymph nodes and tonsils.,


研究领域 (Research Area)

Intestinal immune network for IgA production;Primary immunodeficiency;


注意事项 (Note)

For research use only.

图片 (Image Data)



Western Blot analysis of HeLa, HepG2 cells using AID Polyclonal Antibody..
Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-colon-cancer,
antibody was diluted at 1:100

EnkiLife 优势产品推荐

产品类别	产品货号	产品名称
WB 解决方案	RA10020	2h 极速 WB 即用型全流程试剂盒
	RA10021	4h 快速 WB 即用型全流程试剂盒
	RA10042	5h 畅享版 WB 全流程试剂盒
	RA10037	校准级彩色预染蛋白 Marker (8-180kDa)
	RA10038	校准级彩色预染蛋白 Marker (10-250kDa)
	RA10039	校准级高分子彩色预染蛋白 Marker (25-400kDa)
TSA 多重荧光染色试剂盒	RA10008	TSA 双标三色多重荧光染色试剂盒 (mIHC)
	RA10009	TSA 三标四色多重荧光染色试剂盒 (mIHC)
	RA10010	TSA 四标五色多重荧光染色试剂盒 (mIHC)
	RA10011	TSA 五标六色多重荧光染色试剂盒 (mIHC)
	RA10012	TSA 六标七色多重荧光染色试剂盒 (mIHC)
IHC 检测试剂盒	RA10006	HRP Anti-Mouse/Rabbit IHC Detection System
	RA10007	Polymer-HRP Anti-Mouse/Rabbit IHC Detection System
抗体标记试剂盒	RE80004p	辣根过氧化物酶(HRP)抗体标记试剂盒
	RE80002q	Sulfo-NHS-生物素标记试剂盒
	RE80007p	Cy3 荧光素标记试剂盒
	RE80011p	Fluor488 荧光素标记试剂盒
	RE80017p	Fluor750 荧光素标记试剂盒
	RE80005p	藻红蛋白(R-PE) 抗体快速标记试剂盒
	RE80040	PE-Cy7 串联染料抗体快速标记试剂盒
稳转细胞系构建服务 (免费赠送全膜 WB 验证)	TS-0001	过表达稳转细胞系构建
	TS-0002	敲低稳转细胞系构建
	TS-0003	敲除细胞系构建

限于篇幅，更多产品详见官网：<https://www.enkilife.cn>