产品货号: APRab06838



产品概述 (Summary)

产品名称 (Production Name) Amphiphysin II Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application) WB,IHC,ICC/IF,ELISA

种属反应性 (Reactivity) Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) BIN1

BIN1; AMPHL; Myc box-dependent-interacting protein 1; Amphiphysin II;

别名 (Alternative Names) Amphiphysin-like protein; Box-dependent myc-interacting protein 1; Bridging

integrator 1

基因 ID (Gene ID) 274.0

O00499.Synthesized peptide derived from the C-terminal region of human 蛋白ID (SwissProt ID)

Amphiphysin II.

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000

蛋白分子量 (Molecular Weight) 64kDa

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品货号: APRab06838



研究背景 (Background)

This gene encodes several isoforms of a nucleocytoplasmic adaptor protein, one of which was initially identified as a MYCinteracting protein with features of a tumor suppressor. Isoforms that are expressed in the central nervous system may be involved in synaptic vesicle endocytosis and may interact with dynamin, synaptojanin, endophilin, and clathrin. Isoforms that are expressed in muscle and ubiquitously expressed isoforms localize to the cytoplasm and nucleus and activate a caspase-independent apoptotic process. Studies in mouse suggest that this gene plays an important role in cardiac muscle development. Alternate splicing of the gene results in several transcript variants encoding different isoforms. Aberrant splice variants expressed in tumor cell lines have also been described. [provided by RefSeq, Mar 2016], alternative products:Additional isoforms seem to exist, disease:Defects in BIN1 are the cause of centronuclear myopathy autosomal recessive (ARCNM) [MIM:255200]; also known as autosomal recessive myotubular myopathy. Centronuclear myopathies are congenital muscle disorders characterized by progressive muscular weakness and wasting involving mainly limb girdle, trunk, and neck muscles. It may also affect distal muscles. Weakness may be present during childhood or adolescence or may not become evident until the third decade of life. Ptosis is a frequent clinical feature. The most prominent histopathologic features include high frequency of centrally located nuclei in muscle fibers not secondary to regeneration, radial arrangement of sarcoplasmic strands around the central nuclei, and predominance and hypotrophy of type 1 fibers, function: May be involved in regulation of synaptic vesicle endocytosis. May act as a tumor suppressor and inhibits malignant cell transformation.,PTM:Phosphorylated by protein kinase C.,similarity:Contains 1 BAR domain, similarity: Contains 1 SH3 domain, subunit: Heterodimer with AMPH. Binds SH3GLB1 (By similarity). Binds to SYNJ1 and DNM1 through its SH3 domain, and to clathrin through a region outside of the SH3 domain. Also binds AP2A2. Interacts with the N-terminal transactivation domain of MYC in a manner requiring the integrity of the conserved MYC box regions 1 and 2. Interacts with BIN2. Interacts with HCV NS5A through its SH3 domain., tissue specificity: Ubiquitous. Highest expression in the brain and muscle. Isoform IIA is expressed only in the brain where it is concentrated in axon initial segments and nodes of Ranvier. Isoform BIN1 is widely expressed with highest expression in skeletal muscle.,

研究领域 (Research Area)


Cell Biology

图片 (Image Data)


Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838

产品货号: APRab06838






Western Blot analysis of mouse brain cells using Amphiphysin II Polyclonal Antibody.. Secondary antibody was diluted at



Immunohistochemical analysis of paraffin-embedded human-pancreas, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-pancreas, antibody was diluted at 1:100

产品货号: APRab06838





Immunohistochemical analysis of paraffin-embedded mouse-muscle, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded mouse-muscle, antibody was diluted at 1:100

注意事项 (Note)

For research use only .

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838