

产品名称: AQP3 Rabbit Polyclonal Antibody
产品货号: APRab07071



产品概述 (Summary)

产品名称 (Production Name)	AQP3 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC, IF, ELISA
种属反应性 (Reactivity)	Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	AQP3
别名 (Alternative Names)	AQP3; Aquaporin-3; AQP-3; Aquaglyceroporin-3
基因 ID (Gene ID)	360.0
蛋白 ID (SwissProt ID)	Q92482. The antiserum was produced against synthesized peptide derived from human AQP3. AA range: 165-214

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:100-1:300, ICC/IF 1:50-1:200, ELISA 1:20000-1:40000
蛋白分子量 (Molecular Weight)	

研究背景 (Background)


产品名称: AQP3 Rabbit Polyclonal Antibody
产品货号: AP Rab07071



This gene encodes the water channel protein aquaporin 3. Aquaporins are a family of small integral membrane proteins related to the major intrinsic protein, also known as aquaporin 0. Aquaporin 3 is localized at the basal lateral membranes of collecting duct cells in the kidney. In addition to its water channel function, aquaporin 3 has been found to facilitate the transport of nonionic small solutes such as urea and glycerol, but to a smaller degree. It has been suggested that water channels can be functionally heterogeneous and possess water and solute permeation mechanisms. Alternative splicing of this gene results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Dec 2015], domain: Aquaporins contain two tandem repeats each containing three membrane-spanning domains and a pore-forming loop with the signature motif Asn-Pro-Ala (NPA), function: Forms a water-specific channel that provide kidney medullary collecting duct with high permeability to water, thereby permitting water to move in the direction of an osmotic gradient. May function as a water and urea exit mechanism in antidiuresis in collecting duct cells. Also slightly permeable to urea and glycerol. May play an important role in gastrointestinal tract water transport and in glycerol metabolism, online information: Blood group antigen gene mutation database, polymorphism: AQP3 is responsible for the GIL blood group system. Isoform 2 is detected in GIL-negative individuals that lack functional AQP3, similarity: Belongs to the MIP/aquaporin (TC 1.A.8) family, subcellular location: In collecting ducts of kidney, tissue specificity: Widely expressed in epithelial cells of kidney (collecting ducts) and airways, in keratinocytes, immature dendritic cells and erythrocytes. Isoform 2 is not detectable in erythrocytes at the protein level,

研究领域 (Research Area)

图片 (Image Data)



Immunohistochemistry analysis of AQP3 antibody in paraffin-embedded human lung carcinoma tissue.

注意事项 (Note)

For research use only .