

产品名称: Arginase I Rabbit Polyclonal Antibody
产品货号: AP Rab07111



产品概述 (Summary)

产品名称 (Production Name)	Arginase I Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ARG1
别名 (Alternative Names)	ARG1; Arginase-1; Liver-type arginase; Type I arginase
基因 ID (Gene ID)	383.0
蛋白 ID (SwissProt ID)	P05089.The antiserum was produced against synthesized peptide derived from human ARG1. AA range:61-110

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:10000
蛋白分子量 (Molecular Weight)	35kDa

研究背景 (Background)

产品名称: Arginase I Rabbit Polyclonal Antibody
产品货号: APRab07111




Arginase catalyzes the hydrolysis of arginine to ornithine and urea. At least two isoforms of mammalian arginase exist (types I and II) which differ in their tissue distribution, subcellular localization, immunologic crossreactivity and physiologic function. The type I isoform encoded by this gene, is a cytosolic enzyme and expressed predominantly in the liver as a component of the urea cycle. Inherited deficiency of this enzyme results in argininemia, an autosomal recessive disorder characterized by hyperammonemia. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2011], catalytic activity: L-arginine + H₂O = L-ornithine + urea, cofactor: Binds 2 manganese ions per subunit, disease: Defects in ARG1 are the cause of argininemia (ARGIN) [MIM:207800]; also known as hyperargininemia. Argininemia is a rare autosomal recessive disorder of the urea cycle. Arginine is elevated in the blood and cerebrospinal fluid, and periodic hyperammonemia occurs. Clinical manifestations include developmental delay, seizures, mental retardation, hypotonia, ataxia, progressive spastic quadriplegia, induction: By arginine or homoarginine, online information: Arginase entry, pathway: Nitrogen metabolism; urea cycle; L-ornithine and urea from L-arginine: step 1/1, similarity: Belongs to the arginase family, subunit: Homotrimer,.


研究领域 (Research Area)

Arginine and proline metabolism;

图片 (Image Data)




Western blot analysis of the lysates from HT-29 cells using ARG1 antibody.




Western Blot analysis of 293 cells using Arginase I Polyclonal Antibody


产品名称: Arginase I Rabbit Polyclonal Antibody
产品货号: APRab07111



Immunohistochemical analysis of paraffin-embedded Human liver. 1, Antibody was diluted at 1:200 (4°,overnight) . 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 30min) .



Immunohistochemical analysis of paraffin-embedded Human liver. 1, Antibody was diluted at 1:200 (4°,overnight) . 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 30min) .



Immunohistochemical analysis of paraffin-embedded Human liver. 1, Antibody was diluted at 1:200 (4°,overnight) . 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 30min) .

产品名称: Arginase I Rabbit Polyclonal Antibody
产品货号: APRab07111



注意事项 (Note)

For research use only .