

产品名称: ASCL1 Rabbit Polyclonal Antibody
产品货号: APRab07210



产品概述 (Summary)

产品名称 (Production Name)	ASCL1 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ASCL1
别名 (Alternative Names)	ASCL1; ASH1; BHLHA46; HASH1; Achaete-scute homolog 1; ASH-1; hASH1; Class A basic helix-loop-helix protein 46; bHLHa46
基因 ID (Gene ID)	429.0
蛋白 ID (SwissProt ID)	P50553.The antiserum was produced against synthesized peptide derived from human ASCL1. AA range:109-158

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:20000-1:40000
蛋白分子量 (Molecular Weight)	25kDa

产品名称: ASCL1 Rabbit Polyclonal Antibody
产品货号: APRab07210




研究背景 (Background)


achaete-scute family bHLH transcription factor 1(ASCL1) Homo sapiens This gene encodes a member of the basic helix-loop-helix (BHLH) family of transcription factors. The protein activates transcription by binding to the E box (5'-CANNTG-3'). Dimerization with other BHLH proteins is required for efficient DNA binding. This protein plays a role in the neuronal commitment and differentiation and in the generation of olfactory and autonomic neurons. Mutations in this gene may contribute to the congenital central hypoventilation syndrome (CCHS) phenotype in rare cases. [provided by RefSeq, Jul 2008],function:May play a role at early stages of development of specific neural lineages in most regions of the CNS, and of several lineages in the PNS. Essential for the generation of olfactory and autonomic neurons. Activates transcription by binding to the E box (5'-CANNTG-3'),similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,subunit:Efficient DNA binding requires dimerization with another bHLH protein. Forms a heterodimer with TCF3,

研究领域 (Research Area)

图片 (Image Data)



Immunohistochemistry analysis of ASCL1 antibody in paraffin-embedded human brain tissue.



Western blot analysis of lysate from Jurkat cells, using ASCL1 antibody.

产品名称: ASCL1 Rabbit Polyclonal Antibody
产品货号: APRab07210



注意事项 (Note)

For research use only .