

产品名称: ASIC1 Rabbit Polyclonal Antibody
产品货号: APRab07216



产品概述 (Summary)

产品名称 (Production Name)	ASIC1 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ASIC1 ACCN2 BNAC2
别名 (Alternative Names)	Acid-sensing ion channel 1 (ASIC1;Amiloride-sensitive cation channel 2, neuronal;Brain sodium channel 2;BNAC2)
基因 ID (Gene ID)	41.0
蛋白 ID (SwissProt ID)	P78348.Synthetic peptide from human protein at AA range: 220-280

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:50-1:200,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	70-75kDa

研究背景 (Background)


产品名称: ASIC1 Rabbit Polyclonal Antibody
产品货号: APRab07216



This gene encodes a member of the acid-sensing ion channel (ASIC) family of proteins, which are part of the degenerin/epithelial sodium channel (DEG/ENaC) superfamily. Members of the ASIC family are sensitive to amiloride and function in neurotransmission. The encoded proteins function in learning, pain transduction, touch sensation, and development of memory and fear. Alternatively spliced transcript variants have been described. [provided by RefSeq, Feb 2012], alternative products: The splice variant from ASIC1a described in mouse and rat, which gives rise to an isoform with different N-termini (Asic1b), does not seem to exist in human, function: Cation channel with high affinity for sodium, which is gated by extracellular protons and inhibited by the diuretic amiloride. Also permeable for Ca^{2+} , Li^{+} and K^{+} . Generates a biphasic current with a fast inactivating and a slow sustained phase. Mediates glutamate-independent Ca^{2+} entry into neurons upon acidosis. This Ca^{2+} overloading is toxic for cortical neurons and may be in part responsible for ischemic brain injury. Heteromeric channel assembly seems to modulate channel properties. Functions as a postsynaptic proton receptor that influences intracellular Ca^{2+} concentration and calmodulin-dependent protein kinase II phosphorylation and thereby the density of dendritic spines. Modulates activity in the circuits underlying innate fear., miscellaneous: Potentiated by Ca^{2+} , Mg^{2+} , Ba^{2+} and multivalent cations. Inhibited by anti-inflammatory drugs like salicylic acid (By similarity). Potentiated by FMRFamide-related neuropeptides. PH dependence may be regulated by serine proteases., PTM: Phosphorylation by PKA regulates interaction with PRKCABP and subcellular location. Phosphorylation by PKC may regulate the channel., similarity: Belongs to the amiloride-sensitive sodium channel family., subcellular location: Localizes in synaptosomes at dendritic synapses of neurons. Colocalizes with DLG4., subunit: Homotetramer or heterotetramer with other ASIC proteins (Probable). Interacts with STOM and ACCN1 (By similarity). Interacts with PRKCABP., tissue specificity: Expressed in most or all neurons.,


研究领域 (Research Area)

图片 (Image Data)



Western blot analysis of SW480 lysate, antibody was diluted at 1000. Secondary antibody was diluted at 1:20000

产品名称: ASIC1 Rabbit Polyclonal Antibody
产品货号: APRab07216



Immunohistochemical analysis of paraffin-embedded Human-brain, antibody was diluted at 1:100

注意事项 (Note)

For research use only .