

产品名称: ASIC3 Rabbit Polyclonal Antibody
产品货号: APRab07217



产品概述 (Summary)

产品名称 (Production Name)	ASIC3 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ASIC3 ASIC3; ACCN3; SLNAC1; TNAC1; Acid-sensing ion channel 3; ASIC3; hASIC3;
别名 (Alternative Names)	Amiloride-sensitive cation channel 3; Neuronal amiloride-sensitive cation channel 3; Testis sodium channel 1; hTNaC1
基因 ID (Gene ID)	9311.0
蛋白 ID (SwissProt ID)	Q9UHC3.The antiserum was produced against synthesized peptide derived from the Internal region of human ASIC3. AA range:191-240

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	58kDa

产品名称: ASIC3 Rabbit Polyclonal Antibody
产品货号: APRab07217




研究背景 (Background)

This gene encodes a member of the degenerin/epithelial sodium channel (DEG/ENaC) superfamily. The members of this family are amiloride-sensitive sodium channels that contain intracellular N and C termini, two hydrophobic transmembrane regions, and a large extracellular loop, which has many cysteine residues with conserved spacing. The member encoded by this gene is an acid sensor and may play an important role in the detection of lasting pH changes. In addition, a heteromeric association between this member and acid-sensing (proton-gated) ion channel 2 has been observed as proton-gated channels sensitive to gadolinium. Alternatively spliced transcript variants have been described. [provided by RefSeq, Feb 2012],developmental stage:Expressed in fetal tissues, expression increases in lung and kidney adult tissues.,domain:The PDZ domain-binding motif is involved in interaction with LIN7A, GOPC and MAGI1.,function:Cation channel with high affinity for sodium, which is gated by extracellular protons and inhibited by the diuretic amiloride. Generates a biphasic current with a fast inactivating and a slow sustained phase. In sensory neurons is proposed to mediate the pain induced by acidosis that occurs in ischemic, damaged or inflamed tissue. May be involved in hyperalgesia. May play a role in mechanoreception. Heteromeric channel assembly seems to modulate channel properties.,miscellaneous:Potentiated by FMRFamide-related neuropeptides. Sensitized and potentiated by NPSF. Regulated by lactate and Ca(2+). Inhibited by anti-inflammatory drugs, like salicylic acid (By similarity). Sensitized and potentiated by NPFF.,PTM:Phosphorylated by PKA. Phosphorylated by PKC. In vitro, PRKCABP/PICK-1 is necessary for PKC phosphorylation and activation of a ACCN3/ASIC3-ACCN1/ASIC2b channel, but does not activate a homomeric ACCN3 channel.,similarity:Belongs to the amiloride-sensitive sodium channel family.,subcellular location:Cell surface expression may be stabilized by interaction with LIN7B and cytoplasmic retention by interaction with DLG4. In part cytoplasmic in cochlea cells.,subunit:Homotetramer or heterotetramer with other ASIC proteins (Probable). Interacts with STOM and DLG4 (By similarity). Interacts with LIN7B, MAGI1/BAIAP1, GOPC and ACCN1.,tissue specificity:Expressed by sensory neurons. Strongly expressed in brain, spinal chord, lung, lymph nodes, kidney, pituitary, heart and testis.,

研究领域 (Research Area)

图片 (Image Data)

产品名称: ASIC3 Rabbit Polyclonal Antibody
产品货号: APRab07217



Western blot analysis of lysate from HepG2 cells, using ASIC3 Antibody.



Western Blot analysis of HepG2 cells using ASIC3 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000

注意事项 (Note)

For research use only .