

产品名称: Ataxin-1 Rabbit Polyclonal Antibody
产品货号: AP Rab07251



产品概述 (Summary)

产品名称 (Production Name)	Ataxin-1 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ATXN1
别名 (Alternative Names)	ATXN1; ATX1; SCA1; Ataxin-1; Spinocerebellar ataxia type 1 protein
基因 ID (Gene ID)	6310.0
蛋白 ID (SwissProt ID)	P54253.The antiserum was produced against synthesized peptide derived from human Ataxin 1. AA range:742-791

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:5000-1:10000
蛋白分子量 (Molecular Weight)	87kDa

研究背景 (Background)

产品名称: Ataxin-1 Rabbit Polyclonal Antibody
产品货号: AP Rab07251




ataxin 1(ATXN1) Homo sapiens The autosomal dominant cerebellar ataxias (ADCA) are a heterogeneous group of neurodegenerative disorders characterized by progressive degeneration of the cerebellum, brain stem and spinal cord. Clinically, ADCA has been divided into three groups: ADCA types I-III. ADCAI is genetically heterogeneous, with five genetic loci, designated spinocerebellar ataxia (SCA) 1, 2, 3, 4 and 6, being assigned to five different chromosomes. ADCAII, which always presents with retinal degeneration (SCA7), and ADCAIII often referred to as the 'pure' cerebellar syndrome (SCA5), are most likely homogeneous disorders. Several SCA genes have been cloned and shown to contain CAG repeats in their coding regions. ADCA is caused by the expansion of the CAG repeats, producing an elongated polyglutamine tract in the corresponding protein. The expanded repeats are variable in size and unstable, usually increasing in size when transmitted. At least 2 isoforms are produced. Disease: Defects in ATXN1 are the cause of spinocerebellar ataxia type 1 (SCA1) [MIM:164400]; also known as olivopontocerebellar atrophy I (OPCA I or OPCA1). Spinocerebellar ataxia is a clinically and genetically heterogeneous group of cerebellar disorders. Patients show progressive incoordination of gait and often poor coordination of hands, speech and eye movements, due to cerebellum degeneration with variable involvement of the brainstem and spinal cord. SCA1 belongs to the autosomal dominant cerebellar ataxias type I (ADCA I) which are characterized by cerebellar ataxia in combination with additional clinical features like optic atrophy, ophthalmoplegia, bulbar and extrapyramidal signs, peripheral neuropathy and dementia. SCA1 is caused by expansion of a CAG repeat in the coding region of ATXN1. Longer expansions result in earlier onset and more severe clinical manifestations of the disease. domain: The AXH domain is required for interaction with CIC. function: Binds RNA in vitro. May be involved in RNA metabolism. The expansion of the polyglutamine tract may alter this function. miscellaneous: The self-association seems to be necessary to form nuclear aggregates. online information: Ataxin-1 entry. polymorphism: The poly-Gln region of ATXN1 is highly polymorphic (4 to 39 repeats) in the normal population and is expanded to about 40-83 repeats in spinocerebellar ataxia 1 (SCA1) patients. similarity: Belongs to the ATXN1 family. similarity: Contains 1 AXH domain. subcellular location: Colocalizes with USP7 in the nucleus. subunit: Interacts with CIC (By similarity). Interacts with ANP32A, PQBP1, UBIN, ATXN1L, USP7 and ZNF804A. tissue specificity: Widely expressed throughout the body.

研究领域 (Research Area)


Epigenetics and Nuclear Signaling

图片 (Image Data)


产品名称: Ataxin-1 Rabbit Polyclonal Antibody
产品货号: APRab07251



Immunofluorescence analysis of NIH/3T3 cells, using Ataxin 1 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human lung carcinoma tissue, using Ataxin 1 Antibody. The picture on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using Ataxin-1 Polyclonal Antibody diluted at 1: 500

注意事项 (Note)

For research use only .