

产品名称: Atg16 Rabbit Polyclonal Antibody
产品货号: APRab07285



产品概述 (Summary)

产品名称 (Production Name)	Atg16 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	ATG16L1
别名 (Alternative Names)	ATG16L1; APG16L; Autophagy-related protein 16-1; APG16-like 1
基因 ID (Gene ID)	55054.0
蛋白 ID (SwissProt ID)	Q676U5.Synthesized peptide derived from Atg16 . at AA range: 60-140

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:5000-1:10000
蛋白分子量 (Molecular Weight)	70kDa

研究背景 (Background)

The protein encoded by this gene is part of a large protein complex that is necessary for autophagy, the major process by

产品名称: Atg16 Rabbit Polyclonal Antibody
产品货号: AP Rab07285




which intracellular components are targeted to lysosomes for degradation. Defects in this gene are a cause of susceptibility to inflammatory bowel disease type 10 (IBD10). Several transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Jun 2010],disease:Genetic variations in ATG16L1 are associated with susceptibility to inflammatory bowel disease type 10 (IBD10) [MIM:611081]. IBD is characterized by a chronic relapsing intestinal inflammation. IBD is subdivided into Crohn disease (CD) and ulcerative colitis phenotypes. IBD10 individuals show the phenotype characteristic to CD. It may involve any part of the gastrointestinal tract, but most frequently the terminal ileum and colon. CD is commonly classified as autoimmune disease.,function:Plays an essential role in autophagy.,sequence caution:Wrong choice of CDS.,similarity:Belongs to the WD repeat ATG16 family.,similarity:Contains 7 WD repeats.,subcellular location:Localized to preautophagosomal structure (PAS) where it is involved in the membrane targeting of ATG5.,subunit:Homooligomer. Interacts with ATG5. Part of either the minor and major complexes respectively composed of 4 sets of ATG12-ATG5 and ATG16L1 (400 kDa) or 8 sets of ATG12-ATG5 and ATG16L1 (800 kDa),.

研究领域 (Research Area)

Cell Biology

图片 (Image Data)



Western Blot analysis of Jurkat using Antibody diluted at 1:1000. Secondary antibody was diluted at 1:20000

注意事项 (Note)

For research use only .