

产品名称: Bak Rabbit Polyclonal Antibody
产品货号: APRab07457



产品概述 (Summary)

产品名称 (Production Name)	Bak Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	BAK1
别名 (Alternative Names)	BAK1; BAK; BCL2L7; CDN1; Bcl-2 homologous antagonist/killer; Apoptosis regulator BAK; Bcl-2-like protein 7; Bcl2-L-7
基因 ID (Gene ID)	578.0
蛋白 ID (SwissProt ID)	Q16611.The antiserum was produced against synthesized peptide derived from human Bak. AA range:1-50

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:100-1:300,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	25kDa

产品名称: Bak Rabbit Polyclonal Antibody
产品货号: APRab07457




研究背景 (Background)

The protein encoded by this gene belongs to the BCL2 protein family. BCL2 family members form oligomers or heterodimers and act as anti- or pro-apoptotic regulators that are involved in a wide variety of cellular activities. This protein localizes to mitochondria, and functions to induce apoptosis. It interacts with and accelerates the opening of the mitochondrial voltage-dependent anion channel, which leads to a loss in membrane potential and the release of cytochrome c. This protein also interacts with the tumor suppressor P53 after exposure to cell stress. [provided by RefSeq, Jul 2008],caution:Could be the product of a pseudogene.,domain:Intact BH3 domain is required by BIK, BID, BAK, BAD and BAX for their pro-apoptotic activity and for their interaction with anti-apoptotic members of the Bcl-2 family. Apoptotic members of the Bcl-2 family.,domain:Intact BH3 motif is required by BIK, BID, BAK, BAD and BAX for their pro-apoptotic activity and for their interaction with anti-apoptotic members of the Bcl-2 family.,function:In the presence of an appropriate stimulus, accelerates programmed cell death by binding to, and antagonizing the a repressor Bcl-2 or its adenovirus homolog E1B 19k protein.,function:In the presence of an appropriate stimulus, accelerates programmed cell death by binding to, and antagonizing the a. repressor BCL2 or its adenovirus homolog E1B 19k protein. Low micromolar levels of zinc ions inhibit the promotion of apoptosis.,similarity:Belongs to the Bcl-2 family.,subunit:Forms heterodimers with Bcl-2, E1B 19k protein, and Bcl-X(L).,subunit:Interacts with BCL2A1 (By similarity). Homodimer. Formation of the homodimer is zinc-dependent. Forms heterodimers with BCL2, E1B 19k protein, and BCL2L1 isoform Bcl-X(L).,tissue specificity:Expressed in a wide variety of tissues, with highest levels in the heart and skeletal muscle.,

研究领域 (Research Area)


Cell Biology

图片 (Image Data)




Immunohistochemistry analysis of paraffin-embedded human skeletal muscle tissue, using Bak Antibody. The picture on the right is blocked with the synthesized peptide.


产品名称: Bak Rabbit Polyclonal Antibody
产品货号: APRab07457



Western blot analysis of lysates from 293 cells, using Bak Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using Bak Polyclonal Antibody diluted at 1: 500



Western Blot analysis of 293 cells using Bak Polyclonal Antibody diluted at 1: 500

注意事项 (Note)

For research use only .