产品名称: BAP31 Rabbit Polyclonal Antibody

产品货号: APRab07465



产品概述 (Summary)

产品名称 (Production Name) BAP31 Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit 应用 (Application) WB,ELISA

种属反应性 (Reactivity) Human, Mouse, Rat

产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) BCAP31

BCAP31; BAP31; DXS1357E; B-cell receptor-associated protein 31; BCR-

别名 (Alternative Names) associated protein 31; Bap31; 6C6-AG tumor-associated antigen; Protein CDM;

p28

基因 ID (Gene ID) 10134.0

P51572.The antiserum was produced against synthesized peptide derived 蛋白ID (SwissProt ID)

from human BAP31. AA range:151-200

产品应用 (Application)

稀释比 (Dilution Ratio) WB 1:500-1:2000,ELISA 1:20000-1:40000

蛋白分子量 (Molecular Weight) 28kDa

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: BAP31 Rabbit Polyclonal Antibody

产品货号: APRab07465




研究背景 (Background)

This gene encodes a member of the B-cell receptor associated protein 31 superfamily. The encoded protein is a multi-pass transmembrane protein of the endoplasmic reticulum that is involved in the anterograde transport of membrane proteins from the endoplasmic reticulum to the Golgi and in caspase 8-mediated apoptosis. Microdeletions in this gene are associated with contiguous ABCD1/DXS1375E deletion syndrome (CADDS), a neonatal disorder. Alternative splicing of this gene results in multiple transcript variants. Two related pseudogenes have been identified on chromosome 16. [provided by RefSeq, Jan 2012], disease: Microdeletions in BCAP31 are involved in the contiguous ABCD1/DXS1375E deletion syndrome (CADDS) [MIM:300475]. Patients manifest profound neonatal hypotonia, subsequent failure to thrive, and cholestatic liver disease., function: May play a role in anterograde transport of membrane proteins from the endoplasmic reticulum to the Golgi. May be involved in CASP8-mediated apoptosis., PTM: Cleaved by CASP8 and other caspases., similarity: Belongs to the BCAP29/BCAP31 family., subcellular location: May shuttle between the ER and the intermediate compartment/cis-Golgi complex., subunit: Homodimer and heterodimer with BCAP29. Binds CASP8 (isoform 9) as a complex containing BCAP31, BCAP29, BCL2 and/or BCL2L1. Interacts with VAMP3, VAMP1 and membrane IgD immunoglobulins. May interact with ACTG1 and non-muscle myosin II. Interacts with PTPLB., tissue specificity: Ubiquitous.,

研究领域 (Research Area)

Cell Biology

图片 (Image Data)




Western Blot analysis of various cells using BAP31 Polyclonal Antibody diluted at 1: 2000

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838

产品名称: BAP31 Rabbit Polyclonal Antibody

产品货号: APRab07465





Western Blot analysis of 293T cells using BAP31 Polyclonal Antibody diluted at 1: 2000

注意事项 (Note)

For research use only.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838