

**产品名称: Bcl-2 Rabbit Polyclonal Antibody**  
**产品货号: AP Rab07501**



## 产品概述 (Summary)

|                        |                                  |
|------------------------|----------------------------------|
| 产品名称 (Production Name) | Bcl-2 Rabbit Polyclonal Antibody |
| 描述 (Description)       | Rabbit polyclonal Antibody       |
| 宿主 (Host)              | Rabbit                           |
| 应用 (Application)       | WB,IHC,ICC/IF,ELISA              |
| 种属反应性 (Reactivity)     | Human,Mouse,Rat,Other            |

## 产品性能 (Performance)

|                     |                                                                                                   |
|---------------------|---------------------------------------------------------------------------------------------------|
| 偶联物 (Conjugation)   | Unconjugated                                                                                      |
| 修饰 (Modification)   | Unmodified                                                                                        |
| 同种型 (Isotype)       | IgG                                                                                               |
| 克隆 (Clonality)      | Polyclonal                                                                                        |
| 形式 (Form)           | Liquid                                                                                            |
| 存放说明 (Storage)      | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.          |
| 储存溶液 (Buffer)       | Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N. |
| 纯化方式 (Purification) | Affinity purification                                                                             |

## 免疫原信息 (Immunogen)

|                        |                                                                                                        |
|------------------------|--------------------------------------------------------------------------------------------------------|
| 基因名 (Gene Name)        | BCL2                                                                                                   |
| 别名 (Alternative Names) | BCL2; Apoptosis regulator Bcl-2                                                                        |
| 基因 ID (Gene ID)        | 596.0                                                                                                  |
| 蛋白 ID (SwissProt ID)   | P10415.The antiserum was produced against synthesized peptide derived from human BCL-2. AA range:46-95 |

## 产品应用 (Application)

|                          |                                                                          |
|--------------------------|--------------------------------------------------------------------------|
| 稀释比 (Dilution Ratio)     | WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:200-1:1000,ELISA 1:5000-1:20000 |
| 蛋白分子量 (Molecular Weight) | 26kDa                                                                    |

## 研究背景 (Background)

**产品名称: Bcl-2 Rabbit Polyclonal Antibody**  
**产品货号: AP Rab07501**

---


BCL2, apoptosis regulator(BCL2) Homo sapiens This gene encodes an integral outer mitochondrial membrane protein that blocks the apoptotic death of some cells such as lymphocytes. Constitutive expression of BCL2, such as in the case of translocation of BCL2 to Ig heavy chain locus, is thought to be the cause of follicular lymphoma. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Feb 2016],disease:A chromosomal aberration involving BCL2 may be a cause of follicular lymphoma (FL) [MIM:151430]; also known as type II chronic lymphatic leukemia. Translocation t(14;18)(q32;q21) with immunoglobulin gene regions. BCL2 mutations found in non-Hodgkin lymphomas carrying the chromosomal translocation could be attributed to the Ig somatic hypermutation mechanism resulting in nucleotide transitions.,domain:The BH4 motif is required for anti-apoptotic activity and for interaction with RAF-1.,function:Suppresses apoptosis in a variety of cell systems including factor-dependent lymphohematopoietic and neural cells. Regulates cell death by controlling the mitochondrial membrane permeability. Appears to function in a feedback loop system with caspases. Inhibits caspase activity either by preventing the release of cytochrome c from the mitochondria and/or by binding to the apoptosis-activating factor (APAF-1),online information:Bcl-2 entry,PTM:Phosphorylation/dephosphorylation on Ser-70 regulates anti-apoptotic activity. Growth factor-stimulated phosphorylation on Ser-70 by PKC is required for the anti-apoptosis activity and occurs during the G2/M phase of the cell cycle. In the absence of growth factors, BCL2 appears to be phosphorylated by other protein kinases such as ERKs and stress-activated kinases. Dephosphorylated by protein phosphatase 2A (PP2A),PTM:Proteolytically cleaved by caspases during apoptosis. The cleaved protein, lacking the BH4 motif, has pro-apoptotic activity, causes the release of cytochrome c into the cytosol promoting further caspase activity.,similarity:Belongs to the Bcl-2 family.,subunit:Forms homodimers, and heterodimers with BAX, BAD, BAK and Bcl-X(L). Heterodimerization with BAX requires intact BH1 and BH2 motifs, and is necessary for anti-apoptotic activity (By similarity). Also interacts with APAF1, RAF-1, TP53BP2, BBC3, BCL2L1, MRPL41 and BNIPL. Binding to FKBP8 seems to target BCL2 to the mitochondria and probably interferes with the binding of BCL2 to its targets.,tissue specificity:Expressed in a variety of tissues.,

## 研究领域 (Research Area)


Apoptosis\_Inhibition;Apoptosis\_Mitochondrial;Apoptosis\_Overview;Focal adhesion;Neurotrophin;Amyotrophic lateral sclerosis (ALS);Pathways in cancer;Colorectal cancer;Prostate cancer;Small cell lung cancer;

## 图片 (Image Data)


产品名称: Bcl-2 Rabbit Polyclonal Antibody  
产品货号: APRab07501



Immunofluorescence analysis of HepG2 cells, using BCL-2 Antibody. The picture on the right is blocked with the synthesized peptide.




Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using BCL-2 Antibody. The picture on the right is blocked with the synthesized peptide.




Western blot analysis of lysates from K562 cells, using BCL-2 Antibody. The lane on the right is blocked with the synthesized peptide.


产品名称: Bcl-2 Rabbit Polyclonal Antibody  
产品货号: AP Rab07501



Western Blot analysis of various cells using primary antibody diluted at 1:1000 (4°C overnight) . Secondary antibody: Goat Anti-rabbit IgG IRDye 800 ( diluted at 1:5000, 25°C, 1 hour) . Cell lysate was extracted by Minute™ Plasma Membrane Protein Isolation and Cell Fractionation Kit (SM-005, Inventbiotech,MN,USA) .



Western Blot analysis of various cells using Bcl-2 Polyclonal Antibody diluted at 1: 1000



Western Blot analysis of 3T3 cells using Bcl-2 Polyclonal Antibody diluted at 1: 1000

## 注意事项 (Note)

For research use only .