产品名称: BRCA2 Rabbit Polyclonal Antibody

产品货号: APRab07643



产品概述 (Summary)

产品名称 (Production Name) BRCA2 Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application) IHC,ICC/IF,ELISA

种属反应性 (Reactivity) Human, Rat

产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name) BRCA2

BRCA2; FACD; FANCD1; Breast cancer type 2 susceptibility protein; Fanconi **别名 (Alternative Names)**

anemia group D1 protein

基因 ID (Gene ID) 675.0

P51587.The antiserum was produced against synthesized peptide derived 蛋白ID (SwissProt ID)

from human BRCA2. AA range:31-80

产品应用 (Application)

稀释比 (Dilution Ratio) IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000

蛋白分子量 (Molecular Weight)

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: BRCA2 Rabbit Polyclonal Antibody

产品货号: APRab07643




研究背景 (Background)

Inherited mutations in BRCA1 and this gene, BRCA2, confer increased lifetime risk of developing breast or ovarian cancer. Both BRCA1 and BRCA2 are involved in maintenance of genome stability, specifically the homologous recombination pathway for double-strand DNA repair. The BRCA2 protein contains several copies of a 70 aa motif called the BRC motif, and these motifs mediate binding to the RAD51 recombinase which functions in DNA repair. BRCA2 is considered a tumor suppressor gene, as tumors with BRCA2 mutations generally exhibit loss of heterozygosity (LOH) of the wild-type allele. [provided by RefSeq, Dec 2008], disease: Defects in BRCA2 are a cause of genetic susceptibility to breast cancer (BC) [MIM:612555, 114480]; also called susceptibility to familial breast-ovarian cancer type 2 (BROVCA2). BC is an extremely common malignancy, affecting one in eight women during their lifetime. A positive family history has been identified as major contributor to risk of development of the disease, and this link is striking for early-onset breast cancer. Mutations in BRCA2 are thought to be responsible for some inherited breast cancer. It is linked with male breast cancer., disease: Defects in BRCA2 are the cause of Fanconi anemia complementation group D type 1 (FANCD1) [MIM:605724]. Fanconi anemia [MIM:227650] is an autosomal recessive disorder affecting all bone marrow elements and associated with cardiac, renal, and limb malformations as well as dermal pigmentary changes, function: Involved in double-strand break repair and/or homologous recombination. May participate in S phase checkpoint activation., online information: BRCA2 entry, polymorphism: Genetic variations in BRCA2 may underlie susceptibility to uveal melanoma [MIM:155720]. Uveal melanoma is the most common type of ocular malignant tumor, consisting of overgrowth of uveal melanocytes and often preceded by a uveal nevus.,PTM:Phosphorylated by ATM upon irradiation-induced DNA damage.,similarity:Contains 8 BRCA2 repeats, subunit: Interacts with RAD51 and DSS1. Interacts with ubiquitinated FANCD2. Interacts with PALB2, enables the recombinational repair and checkpoints functions. Interacts with WDR16., tissue specificity: Highest levels of expression in breast and thymus, with slightly lower levels in lung, ovary and spleen.,

研究领域 (Research Area)

Homologous recombination; Pathways in cancer; Pancreatic cancer;

图片 (Image Data)



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using BRCA2 Antibody. The picture on the right is


Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838

产品名称: BRCA2 Rabbit Polyclonal Antibody

产品货号: APRab07643



blocked with the synthesized peptide.



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100 (4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.

注意事项 (Note)

For research use only .

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838