

产品名称: CARKL Rabbit Polyclonal Antibody
产品货号: APRab07931



产品概述 (Summary)

产品名称 (Production Name)	CARKL Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	SHPK
别名 (Alternative Names)	SHPK; CARKL; Sedoheptulokinase; SHK; Carbohydrate kinase-like protein
基因 ID (Gene ID)	23729.0
蛋白 ID (SwissProt ID)	Q9UJH6.The antiserum was produced against synthesized peptide derived from human CARKL. AA range:31-80

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	55kDa


研究背景 (Background)

产品名称: CARKL Rabbit Polyclonal Antibody
产品货号: APRab07931


sedoheptulokinase(SHPK) Homo sapiens The protein encoded by this gene has weak homology to several carbohydrate kinases, a class of proteins involved in the phosphorylation of sugars as they enter a cell, inhibiting return across the cell membrane. Sequence variation between this novel gene and known carbohydrate kinases suggests the possibility of a different substrate, cofactor or changes in kinetic properties distinguishing it from other carbohydrate kinases. The gene resides in a region commonly deleted in cystinosis patients, suggesting a role as a modifier for the cystinosis phenotype. The genomic region is also rich in Alu repetitive sequences, frequently involved in chromosomal rearrangements. [provided by RefSeq, Jul 2008],catalytic activity:ATP + sedoheptulose = ADP + sedoheptulose 7-phosphate.,disease:Deficiency of the SHPK gene in cystinosis patients with a common 57-Kb deletion causes urinary accumulation of sedoheptulose and erythritol.,similarity:Belongs to the FGGY kinase family.,tissue specificity:Strongly expressed in liver, kidney and pancreas. Expressed at lower levels in placenta and heart. Very weakly expressed in lung and brain.,

研究领域 (Research Area)

图片 (Image Data)




Immunohistochemistry analysis of paraffin-embedded human liver carcinoma tissue, using CARKL Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from Jurkat cells, using CARKL Antibody. The lane on the right is blocked with the synthesized peptide.

产品名称: CARKL Rabbit Polyclonal Antibody
产品货号: APRab07931



Western Blot analysis of Jurkat cells using CARKL Polyclonal Antibody

注意事项 (Note)

For research use only .