

产品名称: Cathepsin B Rabbit Polyclonal Antibody  
产品货号: APRab08009



## 产品概述 (Summary)

产品名称 (Production Name)	Cathepsin B Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

## 产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

## 免疫原信息 (Immunogen)

基因名 (Gene Name)	CTSB
别名 (Alternative Names)	CTSB; CPSB; Cathepsin B; APP secretase; APPS; Cathepsin B1
基因 ID (Gene ID)	1508.0
蛋白 ID (SwissProt ID)	P07858.The antiserum was produced against synthesized peptide derived from human Cathepsin B. AA range:168-217

## 产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:5000-1:10000
蛋白分子量 (Molecular Weight)	38,40,25kDa

## 研究背景 (Background)

**产品名称: Cathepsin B Rabbit Polyclonal Antibody**  
**产品货号: APRab08009**




This gene encodes a member of the C1 family of peptidases. Alternative splicing of this gene results in multiple transcript variants. At least one of these variants encodes a preproprotein that is proteolytically processed to generate multiple protein products. These products include the cathepsin B light and heavy chains, which can dimerize to form the double chain form of the enzyme. This enzyme is a lysosomal cysteine protease with both endopeptidase and exopeptidase activity that may play a role in protein turnover. It is also known as amyloid precursor protein secretase and is involved in the proteolytic processing of amyloid precursor protein (APP). Incomplete proteolytic processing of APP has been suggested to be a causative factor in Alzheimer's disease, the most common cause of dementia. Overexpression of the encoded protein has been associated with esophageal adenocarcinoma and other tumors. catalytic activity: Hydrolysis of proteins with broad specificity for peptide bonds. Preferentially cleaves -Arg-Arg-|-Xaa bonds in small molecule substrates (thus differing from cathepsin L). In addition to being an endopeptidase, shows peptidyl-dipeptidase activity, liberating C-terminal dipeptides. function: Thiol protease which is believed to participate in intracellular degradation and turnover of proteins. Has also been implicated in tumor invasion and metastasis. similarity: Belongs to the peptidase C1 family. subcellular location: Identified by mass spectrometry in melanosome fractions from stage I to stage IV. subunit: Dimer of a heavy chain and a light chain cross-linked by a disulfide bond.

### **研究领域 (Research Area)**


Lysosome; Antigen processing and presentation;

### **图片 (Image Data)**




Immunohistochemistry analysis of Cathepsin B antibody in paraffin-embedded human brain tissue.


产品名称: Cathepsin B Rabbit Polyclonal Antibody  
产品货号: APRab08009



Western blot analysis of lysate from COLO cells, using Cathepsin B antibody.




Western Blot analysis of various cells using Cathepsin B Polyclonal Antibody




Immunohistochemical analysis of paraffin-embedded Human colon. 1, Antibody was diluted at 1:200 (4°, overnight) . 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 30min) .

产品名称: Cathepsin B Rabbit Polyclonal Antibody  
产品货号: APRab08009



Immunohistochemical analysis of paraffin-embedded Human colon. 1, Antibody was diluted at 1:200 (4°,overnight) . 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 30min) .



Immunohistochemical analysis of paraffin-embedded Human colon. 1, Antibody was diluted at 1:200 (4°,overnight) . 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 30min) .

### 注意事项 (Note)

For research use only .