

产品名称: Cathepsin E Rabbit Polyclonal Antibody
产品货号: APRab08017



产品概述 (Summary)

产品名称 (Production Name)	Cathepsin E Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Mouse,Rat

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	CTSE
别名 (Alternative Names)	CTSE; Cathepsin E
基因 ID (Gene ID)	1510.0
蛋白 ID (SwissProt ID)	P14091.The antiserum was produced against synthesized peptide derived from human Cathepsin E. AA range:243-292

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	48kDa

研究背景 (Background)

产品名称: Cathepsin E Rabbit Polyclonal Antibody
产品货号: APRab08017




This gene encodes a member of the A1 family of peptidases. Alternative splicing of this gene results in multiple transcript variants. At least one of these variants encodes a preproprotein that is proteolytically processed to generate the mature enzyme. This enzyme, an aspartic endopeptidase, may be involved in antigen processing and the maturation of secretory proteins. Elevated expression of this gene has been observed in neurodegeneration. [provided by RefSeq, Nov 2015],catalytic activity:Similar to cathepsin D, but slightly broader specificity.,function:May have a role in immune function. Probably involved in the processing of antigenic peptides during MHC class II-mediated antigen presentation. May play a role in activation-induced lymphocyte depletion in the thymus, and in neuronal degeneration and glial cell activation in the brain.,PTM:Glycosylated. The nature of the carbohydrate chain varies between cell types. In fibroblasts, the proenzyme contains a high mannose-type oligosaccharide, while the mature enzyme contains a complex-type oligosaccharide. In erythrocyte membranes, both the proenzyme and mature enzyme contain a complex-type oligosaccharide.,PTM:Two forms are produced by autocatalytic cleavage, form I begins at Ile-54, form II begins at Thr-57.,similarity:Belongs to the peptidase A1 family.,subcellular location:The proenzyme is localized to the endoplasmic reticulum and Golgi apparatus, while the mature enzyme is localized to the endosome.,subunit:Homodimer; disulfide-linked.,tissue specificity:Expressed abundantly in the stomach, the Clara cells of the lung and activated B-lymphocytes, and at lower levels in lymph nodes, skin and spleen. Not expressed in resting B-lymphocytes.,

研究领域 (Research Area)


Lysosome;

图片 (Image Data)




Immunohistochemistry analysis of Cathepsin E antibody in paraffin-embedded human lung carcinoma tissue.

产品名称: Cathepsin E Rabbit Polyclonal Antibody
产品货号: APRab08017



Western blot analysis of lysate from HUVEC cells, using Cathepsin E antibody.



Western Blot analysis of various cells using Cathepsin E Polyclonal Antibody

注意事项 (Note)

For research use only .