

产品名称: CD1A/B Rabbit Polyclonal Antibody
产品货号: APRab08259



产品概述 (Summary)

产品名称 (Production Name)	CD1A/B Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC, IF, ELISA
种属反应性 (Reactivity)	Human, Rat, Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	CD1A CD1B
别名 (Alternative Names)	
基因 ID (Gene ID)	909/910
蛋白 ID (SwissProt ID)	P06126/P29016.Synthetic peptide from human protein at AA range: 220-270

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:50-1:200, ICC/IF 1:50-1:200, ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	

研究背景 (Background)

This gene encodes a member of the CD1 family of transmembrane glycoproteins, which are structurally related to the

产品名称: CD1A/B Rabbit Polyclonal Antibody
产品货号: APRab08259




major histocompatibility complex (MHC) proteins and form heterodimers with beta-2-microglobulin. The CD1 proteins mediate the presentation of primarily lipid and glycolipid antigens of self or microbial origin to T cells. The human genome contains five CD1 family genes organized in a cluster on chromosome 1. The CD1 family members are thought to differ in their cellular localization and specificity for particular lipid ligands. The protein encoded by this gene localizes to the plasma membrane and to recycling vesicles of the early endocytic system. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2016],function:Antigen-presenting protein that binds self and non-self lipid and glycolipid antigens and presents them to T-cell receptors on natural killer T-cells.,miscellaneous:During protein synthesis and maturation, CD1 family members bind endogenous lipids that are replaced by lipid or glycolipid antigens when the proteins are internalized and pass through endosomes, before trafficking back to the cell surface.,similarity:Contains 1 Ig-like (immunoglobulin-like) domain.,subcellular location:Subject to intracellular trafficking between the cell membrane and endosomes. Localizes to cell surface lipid rafts.,subunit:Heterodimer with B2M (beta-2-microglobulin). Interacts with CD74.,tissue specificity:Expressed on cortical thymocytes, epidermal Langerhans cells, dendritic cells, on certain T-cell leukemias, and in various other tissues.,


研究领域 (Research Area)

Hematopoietic cell lineage;

图片 (Image Data)



Immunohistochemical analysis of paraffin-embedded Human-tonsil, antibody was diluted at 1:100



产品名称: CD1A/B Rabbit Polyclonal Antibody
产品货号: APRab08259



Immunohistochemical analysis of paraffin-embedded Human-tonsil, antibody was diluted at 1:100

注意事项 (Note)

For research use only .