

产品名称: CD32-C Rabbit Polyclonal Antibody
产品货号: APRab08369



产品概述 (Summary)

产品名称 (Production Name)	CD32-C Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	FCGR2C FCGR2C; CD32; FCG2; IGFR2; Low affinity immunoglobulin gamma Fc region
别名 (Alternative Names)	receptor II-c; IgG Fc receptor II-c; CDw32; Fc-gamma RII-c; Fc-gamma-RIIc; FcRII-c; CD antigen CD32
基因 ID (Gene ID)	9103.0
蛋白 ID (SwissProt ID)	P31995.The antiserum was produced against synthesized peptide derived from human FCGR2C. AA range:251-300

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,ELISA 1:20000-1:40000
蛋白分子量 (Molecular Weight)	35kDa

产品名称: CD32-C Rabbit Polyclonal Antibody
产品货号: APRab08369



研究背景 (Background)

caution:Has sometimes been attributed to correspond to FcR-IIb.,caution:Has sometimes been attributed to correspond to FcR-IIc.,disease:A chromosomal aberration involving FCGR2B is found in a follicular lymphoma. Translocation t(1;22)(q22;q11). The translocation leads to the hyperexpression of the receptor. This may play a role in the tumor progression.,domain:Contains 1 copy of a cytoplasmic motif that is referred to as the immunoreceptor tyrosine-based inhibitor motif (ITIM). This motif is involved in modulation of cellular responses. The phosphorylated ITIM motif can bind the SH2 domain of several SH2-containing phosphatases.,domain:Contains an intracytoplasmic twice repeated motif referred as immunoreceptor tyrosine-based activator motif (ITAM). These motifs are involved in triggering cell activation upon receptors aggregation.,function:Receptor for the Fc region of complexed immunoglobulins gamma. Low affinity receptor. Involved in a variety of effector and regulatory functions such as phagocytosis of immune complexes and modulation of antibody production by B-cells.,function:Receptor for the Fc region of complexed or aggregated immunoglobulins gamma. Low affinity receptor. Involved in a variety of effector and regulatory functions such as phagocytosis of immune complexes and modulation of antibody production by B-cells. Binding to this receptor results in down-modulation of previous state of cell activation triggered via antigen receptors on B-cells (BCR), T-cells (TCR) or via another Fc receptor. Isoform IIB1 fails to mediate endocytosis or phagocytosis. Isoform IIB2 does not trigger phagocytosis.,similarity:Contains 2 Ig-like C2-type (immunoglobulin-like) domains.,subunit:Isoform IIB1 interacts with measles virus N protein. N protein is released in the blood following lysis of measles infected cells. This interaction presumably block inflammatory immune response. Interacts with INPP5D/SHIP1.,tissue specificity:Is the most broadly distributed Fc-gamma-receptor. Expressed in monocyte, neutrophils, macrophages, basophils, eosinophils, Langerhans cells, B-cells, platelets cells and placenta (endothelial cells). Not detected in natural killer cells.,tissue specificity:Isoform IIC1 is detected in monocytes, macrophages, polymorphonuclear cells and natural killer cells.,caution:Has sometimes been attributed to correspond to FcR-IIb.,caution:Has sometimes been attributed to correspond to FcR-IIc.,disease:A chromosomal aberration involving FCGR2B is found in a follicular lymphoma. Translocation t(1;22)(q22;q11). The translocation leads to the hyperexpression of the receptor. This may play a role in the tumor progression.,domain:Contains 1 copy of a cytoplasmic motif that is referred to as the immunoreceptor tyrosine-based inhibitor motif (ITIM). This motif is involved in modulation of cellular responses. The phosphorylated ITIM motif can bind the SH2 domain of several SH2-containing phosphatases.,domain:Contains an intracytoplasmic twice repeated motif referred as immunoreceptor tyrosine-based activator motif (ITAM). These motifs are involved in triggering cell activation upon receptors aggregation.,function:Receptor for the Fc region of complexed immunoglobulins gamma. Low affinity receptor. Involved in a variety of effector and regulatory functions such as phagocytosis of immune complexes and modulation of antibody production by B-cells.,function:Receptor for the Fc region of complexed or aggregated immunoglobulins gamma. Low affinity receptor. Involved in a variety of effector and regulatory functions such as phagocytosis of immune complexes and modulation of antibody production by B-cells. Binding to this receptor results in down-modulation of previous state of cell activation triggered via antigen receptors on B-cells (BCR), T-cells (TCR) or via another Fc receptor. Isoform IIB1 fails to mediate endocytosis or phagocytosis. Isoform IIB2 does not trigger phagocytosis.,similarity:Contains 2 Ig-like C2-type

产品名称: CD32-C Rabbit Polyclonal Antibody
产品货号: APRab08369




(immunoglobulin-like) domains.,subunit:Isoform IIB1 interacts with measles virus N protein. N protein is released in the blood following lysis of measles infected cells. This interaction presumably block inflammatory immune response. Interacts with INPP5D/SHIP1.,tissue specificity:Is the most broadly distributed Fc-gamma-receptor. Expressed in monocyte, neutrophils, macrophages, basophils, eosinophils, Langerhans cells, B-cells, platelets cells and placenta (endothelial cells). Not detected in natural killer cells.,tissue specificity:Isoform IIC1 is detected in monocytes, macrophages, polymorphonuclear cells and natural killer cells.,

研究领域 (Research Area)

B_Cell_Antigen;Fc gamma R-mediated phagocytosis;Systemic lupus erythematosus;

图片 (Image Data)



Western blot analysis of the lysates from HeLa cells using FCGR2C antibody.

注意事项 (Note)

For research use only .