

产品名称: CD71 Rabbit Polyclonal Antibody
产品货号: APRab08446



产品概述 (Summary)

产品名称 (Production Name)	CD71 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	TFRC
别名 (Alternative Names)	TFRC; Transferrin receptor protein 1; TR; TfR; TfR1; Tfr; T9; p90; CD71
基因 ID (Gene ID)	7037.0
蛋白 ID (SwissProt ID)	P02786.The antiserum was produced against synthesized peptide derived from the N-terminal region of human TFRC. AA range:101-150

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:10000-1:20000
蛋白分子量 (Molecular Weight)	85kDa

研究背景 (Background)

产品名称: CD71 Rabbit Polyclonal Antibody
产品货号: AP Rab08446




This gene encodes a cell surface receptor necessary for cellular iron uptake by the process of receptor-mediated endocytosis. This receptor is required for erythropoiesis and neurologic development. Multiple alternatively spliced variants have been identified. [provided by RefSeq, Sep 2015],function:Cellular uptake of iron occurs via receptor-mediated endocytosis of ligand-occupied transferrin receptor into specialized endosomes. Endosomal acidification leads to iron release. The apotransferrin-receptor complex is then recycled to the cell surface with a return to neutral pH and the concomitant loss of affinity of apotransferrin for its receptor. Transferrin receptor is necessary for development of erythrocytes and the nervous system (By similarity). A second ligand, the hereditary hemochromatosis protein HFE, competes for binding with transferrin for an overlapping C-terminal binding site.,induction:Regulated by cellular iron levels through binding of the iron regulatory proteins, IRP1 and IRP2, to iron-responsive elements in the 3'-UTR. Up-regulated upon mitogenic stimulation.,miscellaneous:Canine and feline parvoviruses bind human and feline transferrin receptors and use these receptors to enter and infect cells.,miscellaneous:Serum transferrin receptor (sTfR) is used as a means of detecting erythropoietin (EPO) misuse by athletes and as a diagnostic test for anemia resulting from a number of conditions including rheumatoid arthritis, pregnancy, irritable bowel syndrome and in HIV patients.,PTM:N- and O-glycosylated, phosphorylated and palmitoylated. The serum form is only glycosylated.,PTM:Palmitoylated on both Cys-62 and Cys-67. Cys-62 seems to be the major site of palmitoylation.,PTM:Proteolytically cleaved on Arg-100 to produce the soluble serum form (sTfR).,similarity:Belongs to the peptidase M28 family. M28B subfamily.,similarity:Contains 1 PA (protease associated) domain.,subcellular location:Identified by mass spectrometry in melanosome fractions from stage I to stage IV.,subunit:Homodimer; disulfide-linked. Binds one transferrin or HFE molecule per subunit. Binds the HLA class II histocompatibility antigen, DR1. Interacts with SH3BP3. Interacts with Machupo arenavirus GPC.,

研究领域 (Research Area)


Endocytosis;Hematopoietic cell lineage;

图片 (Image Data)




Western blot analysis of lysate from HUVEC cells, using TFRC Antibody.


产品名称: CD71 Rabbit Polyclonal Antibody
产品货号: APRab08446



Western Blot analysis of HuvEc cells using CD71 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000



Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:100



Immunohistochemical analysis of paraffin-embedded human-lung, antibody was diluted at 1:100

注意事项 (Note)

For research use only .