产品名称: Cdc25C Rabbit Polyclonal Antibody

产品货号: APRab08508



## 产品概述 (Summary)

产品名称 (Production Name) Cdc25C Rabbit Polyclonal Antibody

描述 (Description) Rabbit polyclonal Antibody

宿主 (Host) Rabbit

应用 (Application)WB,IHC,ICC/IF,ELISA种属反应性 (Reactivity)Human,Rat,Mouse

### 产品性能 (Performance)

偶联物 (Conjugation)Unconjugated修饰 (Modification)Unmodified

同种型 (Isotype) IgG

克隆 (Clonality) Polyclonal 形式 (Form) Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid 存放说明 (Storage)

freeze/thaw cycles.

Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% 储存溶液 (Buffer)

New type preservative N.

纯化方式 (Purification) Affinity purification

# 免疫原信息 (Immunogen)

基因名 (Gene Name) CDC25C

CDC25C; M-phase inducer phosphatase 3; Dual specificity phosphatase 别名 (Alternative Names)

Cdc25C

基因 ID (Gene ID) 995.0

P30307.The antiserum was produced against synthesized peptide derived 蛋白ID (SwissProt ID)

from human CDC25C. AA range:183-232

#### 产品应用 (Application)

**稀释比 (Dilution Ratio)** WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:20000-1:40000

蛋白分子量 (Molecular Weight) 53kDa

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技支持) Tel:027-87002838

产品名称: Cdc25C Rabbit Polyclonal Antibody

产品货号: APRab08508




#### 研究背景 (Background)

cell division cycle 25C(CDC25C) Homo sapiens This gene encodes a conserved protein that plays a key role in the regulation of cell division. The encoded protein directs dephosphorylation of cyclin B-bound CDC2 and triggers entry into mitosis. It also suppresses p53-induced growth arrest. Multiple alternatively spliced transcript variants of this gene have been described. [provided by RefSeq, Dec 2015],catalytic activity:Protein tyrosine phosphate + H(2)O = protein tyrosine + phosphate.,developmental stage:Expressed predominantly in G2 phase.,function:Functions as a dosage-dependent inducer in mitotic control. It is a tyrosine protein phosphatase required for progression of the cell cycle. It directly dephosphorylates CDC2 and activate its kinase activity.,PTM:Phosphorylated by CHK1 on Ser-216. This phosphorylation creates a binding site for 14-3-3 protein and inhibits the phosphatase.,similarity:Belongs to the MPI phosphatase family,,similarity:Contains 1 rhodanese domain,,subunit:Interacts with HIV-1 Vpr, thereby inactivating CDC25C phosphatase activity.,


#### 研究领域 (Research Area)

Cell Cycle G1S;Cell Cycle G2M DNA;Oocyte meiosis;Progesterone-mediated oocyte maturation;

### 图片 (Image Data)



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using CDC25C Antibody. The picture on the right is blocked with the synthesized peptide.




Western blot analysis of lysates from HeLa and Jurkat cells, using CDC25C Antibody. The lane on the right is blocked with the synthesized peptide.

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838

产品名称: Cdc25C Rabbit Polyclonal Antibody

产品货号: APRab08508





Western Blot analysis of various cells using Cdc25C Polyclonal Antibody

# 注意事项 (Note)

For research use only .

Web:https://www.enkilife.cn E-mail:order@enkilife.cn (销售) tech@enkilife.cn (技术支持) Tel:027-87002838