

产品名称: CEL Rabbit Polyclonal Antibody
产品货号: APRab08627



产品概述 (Summary)

产品名称 (Production Name)	CEL Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC, IF, ELISA
种属反应性 (Reactivity)	Human, Rat, Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	CEL CEL; BAL; Bile salt-activated lipase; BAL; Bile salt-stimulated lipase; BSSL;
别名 (Alternative Names)	Bucelipase; Carboxyl ester lipase; Cholesterol esterase; Pancreatic lysophospholipase; Sterol esterase
基因 ID (Gene ID)	1056.0
蛋白 ID (SwissProt ID)	P19835.Synthesized peptide derived from the Internal region of human CEL.

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:100-1:300, ICC/IF 1:50-1:200, ELISA 1:20000-1:40000
蛋白分子量 (Molecular Weight)	

产品名称: CEL Rabbit Polyclonal Antibody
产品货号: AP Rab08627




研究背景 (Background)

The protein encoded by this gene is a glycoprotein secreted from the pancreas into the digestive tract and from the lactating mammary gland into human milk. The physiological role of this protein is in cholesterol and lipid-soluble vitamin ester hydrolysis and absorption. This encoded protein promotes large chylomicron production in the intestine. Also its presence in plasma suggests its interactions with cholesterol and oxidized lipoproteins to modulate the progression of atherosclerosis. In pancreatic tumoral cells, this encoded protein is thought to be sequestered within the Golgi compartment and is probably not secreted. This gene contains a variable number of tandem repeat (VNTR) polymorphism in the coding region that may influence the function of the encoded protein. [provided by RefSeq, Jul 2008], catalytic activity: A sterol ester + H₂O = a sterol + a fatty acid., catalytic activity: Triacylglycerol + H₂O = diacylglycerol + a carboxylate., disease: Defects in CEL are a cause of maturity-onset diabetes of the young type 8 with exocrine dysfunction (MODY8) [MIM:609812]; also known as diabetes and pancreatic exocrine dysfunction (DPED). MODY [MIM:606391] is an autosomal dominant form of diabetes mellitus. The pancreas serves both endocrine and exocrine functions. The endocrine cells are found in the islets of Langerhans. They synthesize insulin and other hormones, and are involved in the pathogenesis of diabetes mellitus. The exocrine cells produce bicarbonate and digestive enzymes and are involved in the pathogenesis of pancreatic malabsorption. The localization of the islets within exocrine pancreatic tissue is suggestive of an interdependency and cross-talk between these two cell populations in their normal and in their abnormal function., enzyme regulation: Activated by bile salts containing a 7-hydroxyl group., function: Catalyzes fat and vitamin absorption. Acts in concert with pancreatic lipase and colipase for the complete digestion of dietary triglycerides., similarity: Belongs to the type-B carboxylesterase/lipase family., tissue specificity: Mammary gland and pancreas.,

研究领域 (Research Area)

Steroid biosynthesis; Glycerolipid metabolism;

图片 (Image Data)



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200 (4° overnight) . 2, Tris-EDTA, pH 9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200 (room temperature, 30min) .

产品名称: CEL Rabbit Polyclonal Antibody
产品货号: APRab08627



注意事项 (Note)

For research use only .