

产品名称: CENPF Rabbit Polyclonal Antibody
产品货号: APRab08640



产品概述 (Summary)

产品名称 (Production Name)	CENPF Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	IHC, ICC/IF
种属反应性 (Reactivity)	Human

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	CENPF
别名 (Alternative Names)	CENPF
基因 ID (Gene ID)	1063.0
蛋白 ID (SwissProt ID)	P49454.Synthesized peptide derived from human protein . at AA range: 2190-2270

产品应用 (Application)

稀释比 (Dilution Ratio)	IHC 1:50-1:300, ICC/IF 1:50-1:200
蛋白分子量 (Molecular Weight)	353kDa

研究背景 (Background)

This gene encodes a protein that associates with the centromere-kinetochore complex. The protein is a component of the

产品名称: CENPF Rabbit Polyclonal Antibody
产品货号: APRab08640




nuclear matrix during the G2 phase of interphase. In late G2 the protein associates with the kinetochore and maintains this association through early anaphase. It localizes to the spindle midzone and the intracellular bridge in late anaphase and telophase, respectively, and is thought to be subsequently degraded. The localization of this protein suggests that it may play a role in chromosome segregation during mitosis. It is thought to form either a homodimer or heterodimer.

Autoantibodies against this protein have been found in patients with cancer or graft versus host disease. [provided by RefSeq, Jul 2008],developmental stage:Gradually accumulates during the cell cycle, reaching peak levels in G2 and M phase, and is rapidly degraded upon completion of mitosis.,function:Required for kinetochore function and chromosome segregation in mitosis. Required for kinetochore localization of dynein, LIS1, NDE1 and NDEL1. Regulates recycling of the plasma membrane by acting as a link between recycling vesicles and the microtubule network through its association with STX4 and SNAP25. Acts as a potential inhibitor of pocket protein-mediated cellular processes during development by regulating the activity of RB proteins during cell division and proliferation. May play a regulatory or permissive role in the normal embryonic cardiomyocyte cell cycle and in promoting continued mitosis in transformed, abnormally dividing neonatal cardiomyocytes. Interaction with RB directs embryonic stem cells toward a cardiac lineage. Involved in the regulation of DNA synthesis and hence cell cycle progression, via its C-terminus. Has a potential role regulating skeletal myogenesis and in cell differentiation in embryogenesis. Involved in dendritic cell regulation of T-cell immunity against chlamydia.,PTM:Hyperphosphorylated during mitosis. Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the centromere protein F family.,subcellular location:Relocalizes to the kinetochore/centromere (coronal surface of the outer plate) and the spindle during mitosis. Observed in nucleus during interphase but not in the nucleolus. At metaphase becomes localized to areas including kinetochore and mitotic apparatus as well as cytoplasm. By telophase, is concentrated within the intracellular bridge at either side of the mid-body.,subunit:Interacts with and STX4 (via C-terminus) (By similarity). Interacts (via N-terminus) with RBL1, RBL2 and SNAP25 (By similarity). Self-associates. Interacts with CENP-E and BUBR1 (via C-terminus). Interacts (via C-terminus) with NDE1, NDEL1 and RB1.,

研究领域 (Research Area)

图片 (Image Data)



产品名称: CENPF Rabbit Polyclonal Antibody
产品货号: APRab08640



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200 (4° overnight) .
2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 45min) .

注意事项 (Note)

For research use only .