

产品名称: CEP170 Rabbit Polyclonal Antibody
产品货号: APRab08661



产品概述 (Summary)

产品名称 (Production Name)	CEP170 Rabbit Polyclonal Antibody
描述 (Description)	Rabbit polyclonal Antibody
宿主 (Host)	Rabbit
应用 (Application)	WB,IHC,ICC/IF,ELISA
种属反应性 (Reactivity)	Human,Rat,Mouse

产品性能 (Performance)

偶联物 (Conjugation)	Unconjugated
修饰 (Modification)	Unmodified
同种型 (Isotype)	IgG
克隆 (Clonality)	Polyclonal
形式 (Form)	Liquid
存放说明 (Storage)	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
储存溶液 (Buffer)	Liquid in PBS containing 50% glycerol, 0.5% protective protein and 0.02% New type preservative N.
纯化方式 (Purification)	Affinity purification

免疫原信息 (Immunogen)

基因名 (Gene Name)	CEP170
别名 (Alternative Names)	CEP170; FAM68A; KAB; KIAA0470; Centrosomal protein of 170 kDa; Cep170; KARP-1-binding protein; KARP1-binding protein
基因 ID (Gene ID)	9859.0
蛋白 ID (SwissProt ID)	Q5SW79.The antiserum was produced against synthesized peptide derived from human CEP170. AA range:701-750

产品应用 (Application)

稀释比 (Dilution Ratio)	WB 1:500-1:2000,IHC 1:100-1:300,ICC/IF 1:50-1:200,ELISA 1:20000-1:40000
蛋白分子量 (Molecular Weight)	160kDa

产品名称: CEP170 Rabbit Polyclonal Antibody
产品货号: APRab08661




研究背景 (Background)


centrosomal protein 170(CEP170) Homo sapiens The product of this gene is a component of the centrosome, a non-membraneous organelle that functions as the major microtubule-organizing center in animal cells. During interphase, the encoded protein localizes to the sub-distal appendages of mature centrioles, which are microtubule-based structures thought to help organize centrosomes. During mitosis, the protein associates with spindle microtubules near the centrosomes. The protein interacts with and is phosphorylated by polo-like kinase 1, and functions in maintaining microtubule organization and cell morphology. The human genome contains a putative transcribed pseudogene. Several alternatively spliced transcript variants of this gene have been found, but the full-length nature of some of these variants has not been determined. [provided by RefSeq, Jul 2008],function:Plays a role in microtubule organization.,PTM:Phosphorylated; probably by PLK1. Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the CEP170 family.,similarity:Contains 1 FHA domain.,subcellular location:Associated with the mature mother centriole. Associated with spindle microtubules during mitosis.,subunit:Interacts with PLK1.,

研究领域 (Research Area)

图片 (Image Data)




Immunohistochemistry analysis of paraffin-embedded human testis tissue, using CEP170 Antibody. The picture on the right is blocked with the synthesized peptide.




产品名称: CEP170 Rabbit Polyclonal Antibody
产品货号: APRab08661



Western blot analysis of lysates from HepG2 and Jurkat cells, using CEP170 Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using CEP170 Polyclonal Antibody diluted at 1: 1000



Western Blot analysis of AD293 cells using CEP170 Polyclonal Antibody diluted at 1: 1000

注意事项 (Note)

For research use only .